

Lattice coverings and Delaunay polytopes

Mathieu Dutour Sikirić


Rudjer Bošković Institute, Croatia

January 16, 2014

I. Lattices coverings

Lattice coverings


- ▶ A **lattice** $L \subset \mathbb{R}^n$ is a set of the form $L = \mathbb{Z}v_1 + \cdots + \mathbb{Z}v_n$.
- ▶ A **covering** is a family of balls $B_n(x_i, r)$, $i \in I$ of the same radius r and center x_i such that any $x \in \mathbb{R}^n$ belongs to at least one ball.



- ▶ If L is a lattice, the **lattice covering** is the covering defined by taking the minimal value of $\alpha > 0$ such that $L + B_n(0, \alpha)$ is a covering.

Empty sphere and Delaunay polytopes


- ▶ **Def:** A sphere $S(c, r)$ of center c and radius r in an n -dimensional lattice L is said to be an **empty sphere** if:
 - $\|v - c\| \geq r$ for all $v \in L$,
 - the set $S(c, r) \cap L$ contains $n + 1$ affinely independent points.
- ▶ **Def:** A **Delaunay polytope** P in a lattice L is a polytope, whose vertex-set is $L \cap S(c, r)$.



- ▶ Delaunay polytopes define a tessellation of the Euclidean space \mathbb{R}^n

Lattice covering

- ▶ For a lattice L we define the **covering radius** $\mu(L)$ to be the smallest r such that the family of balls $v + B_n(0, r)$ for $v \in L$ cover \mathbb{R}^n .



- ▶ The covering density has the expression

$$\Theta(L) = \frac{\mu(L)^n \text{vol}(B_n(0, 1))}{\det(L)} \geq 1$$

with

- ▶ $\mu(L)$ being the **largest radius of Delaunay polytopes**
- ▶ or

$$\mu(L) = \max_{x \in \mathbb{R}^n} \min_{y \in L} \|x - y\|$$

Covering minimization and maximization

- ▶ For a given lattice L the only general method for computing $\Theta(L)$ is to compute all Delaunay polytopes.
- ▶ The minimization problem is the problem of **minimizing** $\Theta(L)$ over all lattices L .

The following is known:

- ▶ For $n \leq 5$ the dual root lattice A_n^* is the best lattice covering.
- ▶ For $n = 6$ there is a conjecturally best lattice covering discovered in F. Vallentin PhD thesis.
- ▶ The Leech lattice Λ_{24} is conjectured to be optimal.
- ▶ The function Θ is unbounded from above but we will develop a theory for describing the **local covering maxima**.


The following is known:

- ▶ There is no local covering maxima for $n \leq 5$
- ▶ For $n = 6$ there is exactly one covering maxima: E_6
- ▶ For $n = 7$ there are exactly two covering maxima: E_7 and ER_7 (**Erdahl & Rybnikov** lattice)
- ▶ There is an infinite series DS_n generalizing E_6 and E_7 .

II. Gram matrix formalism

Gram matrix and lattices

- ▶ Denote by S^n the vector space of real symmetric $n \times n$ matrices and $S_{>0}^n$ the convex cone of real symmetric positive definite $n \times n$ matrices.
- ▶ Take a basis (v_1, \dots, v_n) of a lattice L and associate to it the **Gram matrix** $G_v = (\langle v_i, v_j \rangle)_{1 \leq i, j \leq n} \in S_{>0}^n$.
- ▶ Example: take the hexagonal lattice generated by $v_1 = (1, 0)$ and $v_2 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$



$$G_v = \frac{1}{2} \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$$

Isometric lattices

- ▶ Take a basis (v_1, \dots, v_n) of a lattice L with $v_i = (v_{i,1}, \dots, v_{i,n}) \in \mathbb{R}^n$ and write the matrix

$$V = \begin{pmatrix} v_{1,1} & \dots & v_{n,1} \\ \vdots & \ddots & \vdots \\ v_{1,n} & \dots & v_{n,n} \end{pmatrix}$$

and $G_{\mathbf{v}} = V^T V$.

The matrix $G_{\mathbf{v}}$ is defined by $\frac{n(n+1)}{2}$ variables as opposed to n^2 for the basis V .

- ▶ If $M \in S_{>0}^n$, then there exists V such that $M = V^T V$ (Gram Schmidt orthonormalization)
- ▶ If $M = V_1^T V_1 = V_2^T V_2$, then $V_1 = OV_2$ with $O^T O = I_n$ (i.e. O corresponds to an isometry of \mathbb{R}^n).
- ▶ Also if L is a lattice of \mathbb{R}^n with basis \mathbf{v} and u an isometry of \mathbb{R}^n , then $G_{\mathbf{v}} = G_{u(\mathbf{v})}$.

Working with Gram matrices

- ▶ In practice all computations on lattices of \mathbb{R}^n are best done with Gram matrices. For example computing

$$d(x) = \min_{y \in L} \|x - y\|$$

is equivalent to minimizing

$$\min_{y \in \mathbb{Z}^n} (v - y)^T A (x - y)$$

for some $v \in \mathbb{R}^n$ expressed from x .

- ▶ We have the determinant relation

$$\det L = \sqrt{\det G_v}$$

- ▶ In general, Gram matrices are the only information taken into input by programs in lattice theory.
- ▶ They give a parameter space for lattices with a natural topology.

Changing basis

- ▶ If \mathbf{v} and \mathbf{v}' are two basis of a lattice L then $V' = VP$ with $P \in GL_n(\mathbb{Z})$. This implies

$$G_{\mathbf{v}'} = V'^T V' = (VP)^T VP = P^T \{V^T V\} P = P^T G_{\mathbf{v}} P$$

- ▶ If $A, B \in S_{>0}^n$, they are called **arithmetically equivalent** if there is at least one $P \in GL_n(\mathbb{Z})$ such that

$$A = P^T B P$$

- ▶ Lattices up to isometric equivalence correspond to $S_{>0}^n$ up to **arithmetic equivalence**.
- ▶ In practice, **Plesken & Souvignier** wrote a program **isom** for testing arithmetic equivalence and a program **autom** for computing automorphism group of lattices.

III. The lattice covering problem

Equalities and inequalities

- ▶ Take $M = G_v$ with $v = (v_1, \dots, v_n)$ a basis of lattice L .
- ▶ If $V = (w_1, \dots, w_N)$ with $w_i \in \mathbb{Z}^n$ are the vertices of a Delaunay polytope of empty sphere $S(c, r)$ then:

$$\|w_i - c\| = r \text{ i.e. } w_i^T M w_i - 2w_i^T M c + c^T M c = r^2$$

- ▶ Subtracting one obtains


$$\left\{ w_i^T M w_i - w_j^T M w_j \right\} - 2 \left\{ w_i^T - w_j^T \right\} M c = 0$$

- ▶ Inverting matrices, one obtains $M c = \psi(M)$ with ψ linear and so one gets **linear equalities** on M .
- ▶ Similarly $\|w - c\| \geq r$ translates into a **linear inequality** on M : Take $V = (v_0, \dots, v_n)$ a simplex ($v_i \in \mathbb{Z}^n$), $w \in \mathbb{Z}^n$. If one writes $w = \sum_{i=0}^n \lambda_i v_i$ with $1 = \sum_{i=0}^n \lambda_i$, then one has

$$\|w - c\| \geq r \Leftrightarrow w^T M w - \sum_{i=0}^n \lambda_i v_i^T M v_i \geq 0$$

Iso-Delaunay domains

- ▶ Take a lattice L and select a basis v_1, \dots, v_n .
- ▶ We want to assign the Delaunay polytopes of a lattice. Geometrically, this means that



are part of the same iso-Delaunay domain.


- ▶ An iso-Delaunay domain is the assignment of Delaunay polytopes of the lattice.

Primitive iso-Delaunay

- ▶ If one takes a generic matrix M in $S_{>0}^n$, then all its Delaunay are simplices and so no linear equality are implied on M .
- ▶ Hence the corresponding iso-Delaunay domain is of dimension $\frac{n(n+1)}{2}$, they are called **primitive**

Equivalence and enumeration

- ▶ The group $GL_n(\mathbb{Z})$ acts on $S_{>0}^n$ by arithmetic equivalence and preserve the primitive iso-Delaunay domains.
- ▶ Voronoi proved that after this action, there is a finite number of primitive iso-Delaunay domains.
- ▶ **Bistellar flipping** creates one iso-Delaunay from a given iso-Delaunay domain and a facet of the domain. In dim. 2:




- ▶ Enumerating primitive iso-Delaunay domains is done classically:
 - ▶ Find one primitive iso-Delaunay domain.
 - ▶ Find the adjacent ones and reduce by arithmetic equivalence.

The algorithm is graph traversal and iteratively finds all the iso-Delaunay up to equivalence.


The partition of $S_{>0}^2 \subset \mathbb{R}^3$ I

If $q(x, y) = ux^2 + 2vxy + wy^2$ then $q \in S_{>0}^2$ if and only if $v^2 < uw$ and $u > 0$.




The partition of $S_{>0}^2 \subset \mathbb{R}^3$ II

We cut by the plane $u + w = 1$ and get a circle representation.



The partition of $S_{>0}^2 \subset \mathbb{R}^3$ III

Primitive iso-Delaunay domains in $S_{>0}^2$:



IV. SDP optimization

Radius of Delaunay polytope

- ▶ Fix a primitive iso-Delaunay domain, i.e. a collection of simplexes as Delaunay polytopes D_1, \dots, D_m .
- ▶ **Thm:** For every $D_i = \text{Conv}(0, v_1, \dots, v_n)$, the radius of the Delaunay polytope is at most 1 if and only if

$$\begin{pmatrix} 4 & \langle v_1, v_1 \rangle & \langle v_2, v_2 \rangle & \dots & \langle v_n, v_n \rangle \\ \langle v_1, v_1 \rangle & \langle v_1, v_1 \rangle & \langle v_1, v_2 \rangle & \dots & \langle v_1, v_n \rangle \\ \langle v_2, v_2 \rangle & \langle v_2, v_1 \rangle & \langle v_2, v_2 \rangle & \dots & \langle v_2, v_n \rangle \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \langle v_n, v_n \rangle & \langle v_n, v_1 \rangle & \langle v_n, v_2 \rangle & \dots & \langle v_n, v_n \rangle \end{pmatrix} \in \mathcal{S}_{\geq 0}^{n+1}$$

by **Delaunay, Dolbilin, Ryshkov & Shtogrin**.

- ▶ The condition is a semidefinite condition.
- ▶ See for more details
 - ▶ A. Schürmann and F. Vallentin, *Computational approaches to lattice packing and covering problems*, *Discrete & Computational Geometry* **35** (2006) 73–116.
 - ▶ A. Schürmann, *Computational geometry of positive definite quadratic forms*, University Lecture Notes, AMS.

SDP optimization problem

- ▶ Fix a primitive iso-Delaunay domain, i.e. a collection of simplexes as Delaunay polytopes D_1, \dots, D_m .
- ▶ **Thm (Minkowski)**: The function $-\log \det(M)$ is strictly convex on $S_{>0}^n$.
- ▶ Solve the problem
 - ▶ M in the iso-Delaunay domain (linear inequalities),
 - ▶ the Delaunay D_i have radius at most 1 (semidefinite condition),
 - ▶ **minimize** $-\log \det(M)$ (strictly convex).
- ▶ **Thm**: Given an iso-Delaunay domain LT , there exist a **unique** lattice, which minimize the covering density over LT .
- ▶ The above problem is solved by the **interior point methods** implemented in **MAXDET** by **Vandenbergh, Boyd & Wu**. Unicity comes from the strict convexity of the objective function.

Solving the minimum covering problem

- ▶ The lattice covering problem is to find a lattice covering of minimal density.
- ▶ The solution of the SDP problem by interior point methods does not give exact solutions but approximate solutions available at any precision.
- ▶ The exact solution is expressible with algebraic integers once one knows which inequations are satisfied with equality.
- ▶ The method for solving the lattice covering problem in dimension n is thus:
 - ▶ Enumerate all iso-Delaunay domains LT up to equivalence
 - ▶ solve the SDP on all the domains
 - ▶ Take the one(s) of minimum covering density
- ▶ **Pb:** 222 primitive iso-Delaunay domains in dimension 5 (**Baranovski, Ryshkov, Engel & Grishukhin**) and at least 200 millions in dimension 6 (**Engel**).
This is not practical at all

V. iso-Delaunay
domains of
 $S_{>0}^n$ -spaces


$S_{>0}^n$ -spaces

- ▶ A $S_{>0}^n$ -space is a vector space \mathcal{SP} of S^n , which intersect $S_{>0}^n$.
- ▶ We want to describe the Delaunay decomposition of matrices $M \in S_{>0}^n \cap \mathcal{SP}$.
- ▶ Motivations:
 - ▶ The enumeration of iso-Delaunay is done up to dimension 5 but certainly not for higher dimension.
 - ▶ We hope to find some good **covering** by selecting judicious \mathcal{SP} . This is a search for best but unproven to be optimal coverings.
- ▶ A iso-Delaunay in \mathcal{SP} is an open convex polyhedral set included in $S_{>0}^n \cap \mathcal{SP}$, for which every element has the **same Delaunay decomposition**.
- ▶ Typical choice of a space \mathcal{SP} are the space of forms invariant under a finite integral matrix group G . In that case finiteness of the set of iso-Delaunay up to equivalence is proved.
- ▶ Dimension of the space \mathcal{SP} is typically no larger than 4.

Lifted Delaunay decomposition

- ▶ The Delaunay polytopes of a lattice L correspond to the facets of the convex cone $\mathcal{C}(L)$ with vertex-set:


$$\{(x, \|x\|^2) \text{ with } x \in L\} \subset \mathbb{R}^{n+1}.$$



- ▶ H. Edelsbrunner, N.R. Shah, *Incremental Topological Flipping Works for Regular Triangulations*, *Algorithmica* **15** (1996) 223–241.

Generalized bistellar flips

- ▶ The “glued” Delaunay form a Delaunay decomposition for a matrix M in the (\mathcal{SP}, L) -iso-Delaunay satisfying to $f(M) = 0$.
- ▶ The flipping break those Delaunays in a different way.
- ▶ Two triangulations of \mathbb{Z}^2 correspond in the lifting to:



- ▶ The polytope represented is called the **repartitioning polytope**. It has two partitions into Delaunay polytopes.
- ▶ The lower facets correspond to one tessellation, the upper facets to the other tessellation.

Enumeration technique

- ▶ Find a primitive (\mathcal{SP}, L) -iso-Delaunay domain, insert it to the list as undone.
- ▶ Iterate
 - ▶ For every undone primitive (\mathcal{SP}, L) -iso-Delaunay domain, compute the facets.
 - ▶ Eliminate **redundant** inequalities.
 - ▶ For every **non-redundant** inequality realize the flipping, i.e. compute the adjacent primitive (\mathcal{SP}, L) -iso-Delaunay domain. If it is new, then add to the list as undone.
- ▶ See for full details
 - ▶ M. Dutour Sikirić, F. Vallentin and A. Schürmann, *A generalization of Voronoi's reduction theory and applications*, Duke Math. J. 142 (2008), 127–164.

Best known lattice coverings

d	lattice / covering density Θ		
1	\mathbb{Z}^1 1	13	L_{13}^c (DSV) 7.762108
2	A_2^* (Kershner) 1.209199	14	L_{14}^c (DSV) 8.825210
3	A_3^* (Bambah) 1.463505	15	L_{15}^c (DSV) 11.004951
4	A_4^* (Delaunay & Ryshkov) 1.765529	16	A_{16}^* (DSV) 15.310927
5	A_5^* (Ryshkov & Baranovski) 2.124286	17	A_{17}^9 (DSV) 12.357468
6	L_6^c (Vallentin) 2.464801	18	A_{18}^* 21.840949
7	L_7^c (Schürmann & Vallentin) 2.900024	19	A_{19}^{10} (DSV) 21.229200
8	L_8^c (Schürmann & Vallentin) 3.142202	20	A_{20}^7 (DSV) 20.366828
9	L_9^c (DSV) 4.268575	21	A_{21}^{11} (DSV) 27.773140
10	L_{10}^c (DSV) 5.154463	22	Λ_{22}^* (Smith) \leq 27.8839
11	L_{11}^c (DSV) 5.505591	23	Λ_{23}^* (Smith, MDS) 15.3218
12	L_{12}^c (DSV) 7.465518	24	Leech 7.903536

- ▶ For $n \leq 5$ the results are definitive.
- ▶ The lattices A_n^r for r dividing $n + 1$ are the Coxeter lattices. They are often good coverings and they are used for perturbations.
- ▶ For dimensions 10 and 12 we use laminations over Coxeter lattices of dimension 9 and 11.
- ▶ Leech lattice is conjecturally optimal (it is local optimal Schürmann & Vallentin)

VI. Quadratic functions and the Erdahl cone

The Erdahl cone

- ▶ Denote by $E_2(n)$ the vector space of degree 2 polynomial functions on \mathbb{R}^n . We write $f \in E_2(n)$ in the form

$$f(x) = a_f + b_f \cdot x + Q_f[x]$$

with $a_f \in \mathbb{R}$, $b_f \in \mathbb{R}^n$ and Q_f a $n \times n$ symmetric matrix

- ▶ The Erdahl cone is defined as

$$\text{Erdahl}(n) = \{f \in E_2(n) \text{ such that } f(x) \geq 0 \text{ for } x \in \mathbb{Z}^n\}$$

- ▶ It is a convex cone, which is non-polyhedral since defined by an infinity of inequalities.
- ▶ The group acting on $\text{Erdahl}(n)$ is $\text{AGL}_n(\mathbb{Z})$, i.e. the group of affine integral transformations

$$x \mapsto b + Px \text{ for } b \in \mathbb{Z}^n \text{ and } P \in \text{GL}_n(\mathbb{Z})$$

Scalar product

- ▶ **Def:** If $f, g \in E_2(n)$, then:

$$\langle f, g \rangle = a_f a_g + \langle b_f, b_g \rangle + \langle Q_f, Q_g \rangle$$

- ▶ **Def:** For $v \in \mathbb{Z}^n$, define $ev_v(x) = (1 + v \cdot x)^2$.
- ▶ We have

$$\langle f, ev_v \rangle = f(v)$$

- ▶ Thus finding the rays of $Erdahl(n)$ is a dual description problem with an infinity of inequalities and infinite group acting on it.
- ▶ If $f \in Erdahl(n)$ then Q_f is positive semidefinite.
- ▶ **Def:** We also define

$$Erdahl_{>0}(n) = \{f \in Erdahl(n) : Q_f \text{ positive definite}\}$$

Relation with Delaunay polytope

- ▶ If D is a Delaunay polytope of a lattice $L = \mathbb{Z}v_1 + \cdots + \mathbb{Z}v_n$ of empty sphere $S(c, r)$ then we define the function

$$f_{D, \mathbf{v}} : \mathbb{Z}^n \rightarrow \mathbb{R}$$
$$x = (x_1, \dots, x_n) \mapsto \left\| \sum_{i=1}^n x_i v_i - c \right\|^2 - r^2$$

Clearly $f_{D, \mathbf{v}} \in \text{Erdahl}_{>0}(n)$.

- ▶ The **perfection rank** of a Delaunay polytope is the dimension of the face it defines in $\text{Erdahl}(n)$.
- ▶ **Def:** If $f \in \text{Erdahl}(n)$ then

$$Z(f) = \{v \in \mathbb{Z}^n : f(v) = 0\}$$

- ▶ **Thm:** If $f \in \text{Erdahl}(n)$ then there exist a lattice L_f and a lattice L' containing a Delaunay polytope D_f such that

$$Z(f) = D_f + L_f$$

- ▶ We have $\dim L' + \dim L_f \leq n$. In case of equality $Z(f)$ is called a **Delaunay polyhedra**.

Perfect Delaunay polytopes/polyhedra

- ▶ **Def:** If D is a n -dimensional Delaunay polyhedra then we define

$$\text{Dom}_{\mathbf{v}} D = \sum_{\mathbf{v} \in D} \mathbb{R}_+ e_{\mathbf{v}}$$

- ▶ We have $\langle f_{D, \mathbf{v}}, \text{Dom}_{\mathbf{v}} D \rangle = 0$.
- ▶ **Def:** D is **perfect** if $\text{Dom } D$ is of dimension $\binom{n+2}{2} - 1$ that is if the perfection rank is 1.
- ▶ This implies that f_D generates an extreme ray of $\text{Erdahl}(n)$ and f_D is rational.
- ▶ A perfect n -dimensional Delaunay polytope has at least $\binom{n+2}{2} - 1$ vertices. There is only one way to embed it as a Delaunay polytope of a lattice.
- ▶ Perfect Delaunay polytopes are remarkable and rare objects.

VII. Covering maxima, pessima and their characterization

Eutacticity

- ▶ If $f \in \text{Erdahl}_{>0}(n)$ then define μ_f and c_f such that

$$f(x) = Q_f[x - c_f] - \mu_f$$

Then define

$$u_f(x) = (1 + c_f \cdot x)^2 + \frac{\mu_f}{n} Q_f^{-1}[x]$$

- ▶ **Def:** $f \in \text{Erdahl}_{>0}(n)$ is **eutactic** if u_f is in the relative interior of $\text{Dom } f$.
- ▶ **Def:** Take a Delaunay polytope P for a quadratic form Q of center c_P and square radius μ_P . P is called **eutactic** if there are $\alpha_v > 0$ so that

$$\left\{ \begin{array}{l} 1 = \sum_{v \in \text{vert } P} \alpha_v, \\ 0 = \sum_{v \in \text{vert } P} \alpha_v (v - c_P), \\ \frac{\mu_P}{n} Q^{-1} = \sum_{v \in \text{vert } P} \alpha_v (v - c_P)(v - c_P)^T. \end{array} \right.$$

Covering maxima

- ▶ A given lattice L is called a **covering maxima** if for any lattice L' near L we have $\Theta(L') < \Theta(L)$.
- ▶ **Thm**: For a lattice L the following are equivalent:
 - ▶ L is a covering maxima
 - ▶ Every Delaunay polytope of maximal circumradius of L is perfect and eutactic.
- ▶ The following are covering maxima:

name	# vertices	# orbits Delaunay polytopes
E_6	27	1
E_7	56	2
ER_7	35	4
O_{10}	160	6
BW_{16}	512	4
O_{23}	94208	5
Λ_{23}	47104	709

- ▶ **Thm**: For any $n \geq 6$ there exist one lattice $L(DS_n)$ which is a covering maxima.
There is only one perfect Delaunay polytope $P(DS_n)$ of maximal radius in $L(DS_n)$.

The infinite series

- ▶ For n even $P(DS_n)$ is defined as the lamination over D_{n-1} of
 - ▶ one vertex
 - ▶ the half cube $\frac{1}{2}H_{n-1}$
 - ▶ the cross polytope CP_{n-1}

For $n = 6$, it is E_6 .

- ▶ For n odd as the lamination over D_{n-1} of
 - ▶ the cross polytope CP_{n-1}
 - ▶ the half cube $\frac{1}{2}H_{n-1}$
 - ▶ the cross polytope CP_{n-1}

For $n = 7$, it is E_7 .

- ▶ **Conj:** The lattice DS_n has the following properties:
 - ▶ $L(DS_n)$ has the maximum covering density among all n -dim. covering maxima
 - ▶ Among all perfect Delaunay polytopes, $P(DS_n)$ has
 - ▶ maximum number of vertices
 - ▶ maximum volume

If true this would imply Minkowski conjecture by results of

- ▶ U. Shapira and B. Weiss, *Stable Lattices and the Diagonal Group*, preprint

Pessimism and Morse function property

- ▶ For a lattice L let us denote $D_{crit}(L)$ the space of direction d of deformation of L such that Θ increases in the direction d .
- ▶ **Def:** A lattice L is said to be a covering **pessimism** if the space D_{crit} is of measures 0.
- ▶ **Thm:** If the Delaunay polytopes of maximum circumradius of a lattice L are eutactic and are not simplices then L is a pessimism.

name	# vertices	# orbits Delaunay polytopes
\mathbb{Z}^n	2^n	1
D_4	8	1
D_n ($n \geq 5$)	2^{n-1}	2
E_6^*	9	1
E_7^*	16	1
E_8	16	2
K_{12}	81	4

- ▶ **Thm:** The covering density function $Q \mapsto \Theta(Q)$ is a topological Morse function if and only if $n \leq 3$.

VIII. Enumeration of perfect Delaunay polytopes

Perfect Delaunay polytope

- ▶ There is a finite number of them in each dimension n . Known results:

dim.	perfect Delaunay	authors
1	$[0, 1]$ in \mathbb{Z}	
2	\emptyset	
3	\emptyset	
4	\emptyset	
5	\emptyset	↑ (Deza, Laurent & Grishukhin)
6	2_{21} in E_6	(Deza & Dutour)
7	3_{21} in E_7 and ER_7 in $L(ER_7)$	(Dutour Sikirić)
8	≥ 27	(Dutour Sikirić & Rybnikov)
9	≥ 100000	(Dutour Sikirić)

- ▶ **Thm:** There exist perfect Delaunay polytopes D such that $\mathbb{Z}D \neq \mathbb{Z}^n$ (dimension $n \geq 13$, Rybnikov & Dutour Sikirić).
- ▶ **Thm:** There exist lattices with several perfect Delaunay polytopes (dimension 15 and 23, Rybnikov & Dutour Sikirić).
- ▶ **Thm:** For $n \geq 6$ there exist a perfect Delaunay polytope with exactly $\binom{n+2}{2} - 1$ vertices (Erdahl & Rybnikov) ER_n .

Extreme rays of $Erdahl(n)$

- ▶ **Def:** If $f \in Erdahl_{>0}(n)$ then we define

$$\text{Dom } f = \sum_{v \in Z(f)} \mathbb{R}_+ e_v$$

- ▶ We have $\langle f, \text{Dom } f \rangle = 0$.
- ▶ **Thm (Erdahl):** The extreme rays of $Erdahl(n)$ are:

(a) The constant function 1.

(b) The functions

$$(a_1 x_1 + \cdots + a_n x_n + \beta)^2$$

with (a_1, \dots, a_n) not collinear to an integral vector.

(c) The functions f such that $Z(f)$ is a perfect Delaunay polyhedra.

- ▶ Note that if $f \in Erdahl(n)$ with $Z(f)$ a Delaunay polyhedra, then there exist a lattice L' of dimension $k \leq n$, a Delaunay polytope D of L' , a basis \mathbf{v}' of L' and a function $\phi \in \text{AGL}_n(\mathbb{Z})$ such that

$$f \circ \phi(x_1, \dots, x_n) = f_{D, \mathbf{v}'}(x_1, \dots, x_k)$$

Delaunay polyhedra retract

- ▶ For a function $f \in \text{Erdahl}(n)$ a **proper decomposition** is a pair (g, h) with $f = g + h$, $g \in \text{Erdahl}(n)$ and $h(x) \geq 0$ for $x \in \mathbb{R}^n$.

- ▶ **Lem:** For a proper decomposition we have

$$\text{Vect } Z(f) + \text{Ker } Q_f \subset \text{Ker } Q_h$$

and there exist a proper decomposition with equality.

- ▶ Fix an integral complement L' of $\text{Vect } Z(f) + \text{Ker } Q_f$. A proper decomposition is called **extremal** if $\det Q_h|_{L'}$ is maximal among all proper decompositions.
- ▶ **Thm:** For $f \in \text{Erdahl}(n)$, there exist a unique extremal decomposition. For it we have that $Z(g)$ is a Delaunay polyhedra.
- ▶ **Conj:** The decomposition depends continuously on $f \in \text{Erdahl}(n)$.
- ▶ On the other hand in a neighborhood of $f \in \text{Erdahl}(n)$ we can have an infinity of Delaunay polyhedra.

Enumeration of perfect Delaunay polyhedra

- ▶ From a given n -dimensional Delaunay polyhedron P of form f we can define the local cone

$$\text{Loc}(f) = \{g \in E_2(n) \text{ s.t. } g(x) \geq 0 \text{ for } x \in Z(f)\}.$$

We set to define the degeneracy $d(P) = \dim L_f$.

- ▶ **Thm:** For a Delaunay polyhedron P let $(P_i)_{i \in I}$ the set of Delaunay polyhedra of degeneracy $d(P) - 1$ and perfection rank $r(P) - 1$. P_i and P_j are adjacent if $P_i \cap P_j$ is of perfection rank $r(P) - 2$. The obtained graph is connected.
- ▶ **Thm:** In a fixed dimension n there exist an algorithm for enumerating the perfect Delaunay polytopes of dimension n . The algorithm is iterative. It relies on dual description. If the degeneracy rank is $d > 0$ then we find a sub Delaunay polyhedron of degeneracy $d - 1$, finds its facets and do the liftings. This requires knowing the facets of CUT_{n+1} .
- ▶ **Thm:** In dimension 7 there is only 3_{21} and ER_7 .