

“Degree Spectra on a Cone” for Polish Spaces

Takayuki Kihara

Japan Advanced Institute of Science and Technology (JAIST)

Joint Work with

Arno Pauly (University of Cambridge, UK)

Sets and Computations, IMS, Singapore, 2015

Let $\mathcal{B}_\alpha^*(\mathbf{X})$ be the Banach algebra of bounded real valued Baire class α functions on \mathbf{X} w.r.t. the supremum norm and pointwise operation.

Main Problem (Motto Ros)

Suppose that \mathbf{X} is an uncountable Polish space.

Is the Banach algebra $\mathcal{B}_n^*(\mathbf{X})$ linearly isometric (ring isomorphic) to either $\mathcal{B}_n^*(\mathbb{R})$ or $\mathcal{B}_n^*(\mathbb{R}^{\aleph})$ for some $n \in \omega$?

Let $\mathcal{B}_\alpha^*(\mathbf{X})$ be the Banach algebra of bounded real valued Baire class α functions on \mathbf{X} w.r.t. the supremum norm and pointwise operation.

Main Problem (Motto Ros)

Suppose that \mathbf{X} is an uncountable Polish space.

Is the Banach algebra $\mathcal{B}_n^*(\mathbf{X})$ linearly isometric (ring isomorphic) to either $\mathcal{B}_n^*(\mathbb{R})$ or $\mathcal{B}_n^*(\mathbb{R}^{\mathbb{N}})$ for some $n \in \omega$?

- We apply [Recursion Theory](#) (a.k.a. Computability Theory) to solve Motto Ros' problem!
- More specifically, an invariant which we call [degree co-spectrum](#), a collection of Turing ideals realized as lower Turing cones of points of a Polish space, plays a key role.
- The key idea is measuring the quantity of all possible [Scott ideals](#) (ω -models of [WKL₀](#)) realized within the degree co-spectrum (on a cone) of a given space.

Background in Banach Space Theory

- The basic theory on the Banach spaces $\mathcal{B}_\alpha(\mathbf{X})$ has been studied by Bade, Dachiell, Jayne and others in 1970s.
- Jayne (1974) proved an analogue of the *Banach-Stone Theorem* and the *Gel'fand-Kolmogorov Theorem* for Baire classes, that is, the α -th level Baire structure of a space \mathbf{X} is determined by the ring structure of the Banach algebra $\mathcal{B}_\alpha^*(\mathbf{X})$, and vice versa.

Background in Banach Space Theory

- The basic theory on the Banach spaces $\mathcal{B}_\alpha(\mathbf{X})$ has been studied by Bade, Dachiell, Jayne and others in 1970s.
- Jayne (1974) proved an analogue of the *Banach-Stone Theorem* and the *Gel'fand-Kolmogorov Theorem* for Baire classes, that is, the α -th level Baire structure of a space \mathbf{X} is determined by the ring structure of the Banach algebra $\mathcal{B}_\alpha^*(\mathbf{X})$, and vice versa.

(Jayne) An α -th level Baire isomorphism is a bijection $f : \mathbf{X} \rightarrow \mathbf{Y}$ s.t. $E \subseteq \mathbf{X}$ is of additive Baire class α iff $f[E] \subseteq \mathbf{Y}$ is of additive Baire class α .

Theorem (Jayne 1974)

The following are equivalent for a realcompact space \mathbf{X} :

- 1 \mathbf{X} is α -th level Baire isomorphic to \mathbf{Y} .
- 2 $\mathcal{B}_\alpha^*(\mathbf{X})$ is linearly isometric to $\mathcal{B}_\alpha^*(\mathbf{Y})$.
- 3 $\mathcal{B}_\alpha^*(\mathbf{X})$ is ring isomorphic to $\mathcal{B}_\alpha^*(\mathbf{Y})$.

Recall that Baire classes and Borel classes coincide in separable metrizable spaces (Lebesgue-Hausdorff).

An *n*-th level Borel isomorphism is a bijection $f : X \rightarrow Y$ s.t.

$$E \subseteq X \text{ is } \Sigma_{\sim n+1}^0 \iff f[E] \subseteq Y \text{ is } \Sigma_{\sim n+1}^0.$$

By Jayne's theorem (1974), Motto Ros' problem is reformulated as:

Recall that Baire classes and Borel classes coincide in separable metrizable spaces (Lebesgue-Hausdorff).

An *n-th level Borel isomorphism* is a bijection $f : X \rightarrow Y$ s.t.

$$E \subseteq X \text{ is } \Sigma_{\sim n+1}^0 \iff f[E] \subseteq Y \text{ is } \Sigma_{\sim n+1}^0.$$

By Jayne's theorem (1974), Motto Ros' problem is reformulated as:

The Second-Level Borel Isomorphism Problem

Find an uncountable Polish space which is second-level Borel isomorphic neither to \mathbb{R} nor to $\mathbb{R}^{\mathbb{N}}$.

Consequently, Motto Ros' problem is the problem on the [second level](#) Borel isomorphic classification of Polish spaces.

Consequently, Motto Ros' problem is the problem on the **second level** Borel isomorphic classification of Polish spaces.

*“We show that any two uncountable Polish spaces that are countable unions of sets of finite dimension are Borel isomorphic at the **second level**, and consequently at all **higher levels**. Thus the **first level** and **zero-th level** (i.e. homeomorphisms) appear to be the **only** levels giving rise to nontrivial classifications of Polish spaces.”*

J. E. Jayne and C. A. Rogers, Borel isomorphisms at the first level I, *Mathematika* **26** (1979), 125-156.

Consequently, Motto Ros' problem is the problem on the **second level** Borel isomorphic classification of Polish spaces.

*"We show that any two uncountable Polish spaces that are countable unions of sets of finite dimension are Borel isomorphic at the **second level**, and consequently at all **higher levels**. Thus the **first level** and **zero-th level** (i.e. homeomorphisms) appear to be the **only** levels giving rise to nontrivial classifications of Polish spaces."*

J. E. Jayne and C. A. Rogers, Borel isomorphisms at the first level I, *Mathematika* **26** (1979), 125-156.

- At that time, almost no nontrivial proper infinite dimensional Polish spaces had been discovered yet.
- Perhaps, it had been expected that the structure of proper infinite dim. Polish spaces is simple

Consequently, Motto Ros' problem is the problem on the **second level** Borel isomorphic classification of Polish spaces.

*"We show that any two uncountable Polish spaces that are countable unions of sets of finite dimension are Borel isomorphic at the **second level**, and consequently at all **higher levels**. Thus the **first level** and **zero-th level** (i.e. homeomorphisms) appear to be the **only levels** giving rise to nontrivial classifications of Polish spaces."*

J. E. Jayne and C. A. Rogers, Borel isomorphisms at the first level I, *Mathematika* **26** (1979), 125-156.

- At that time, almost no nontrivial proper infinite dimensional Polish spaces had been discovered yet.
- Perhaps, it had been expected that the structure of proper infinite dim. Polish spaces is simple — this conclusion was too hasty!
- By using **Recursion Theory**, we reveal that the **second level** Borel isomorphic classification of Polish spaces is highly nontrivial!

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$, i.e., X_α is **compact metrizable**, and if $X_\alpha \setminus \mathbf{C} = U_1 \sqcup U_2$ for some nonempty open U_1, U_2 , then \mathbf{C} must be infinite dimensional.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$, i.e., X_α is **compact metrizable**, and if $X_\alpha \setminus \mathbf{C} = U_1 \sqcup U_2$ for some nonempty open U_1, U_2 , then \mathbf{C} must be infinite dimensional.
- 2 X_α possesses Haver's property \mathbf{C} (hence, **weakly infinite dimensional**) for any $\alpha < 2^{\aleph_0}$.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$, i.e., X_α is compact metrizable, and if $X_\alpha \setminus C = U_1 \sqcup U_2$ for some nonempty open U_1, U_2 , then C must be infinite dimensional.
- 2 X_α possesses Haver's property C (hence, weakly infinite dimensional) for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is not n -th level Borel isomorphic to X_β .

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$, i.e., X_α is **compact metrizable**, and if $X_\alpha \setminus C = U_1 \sqcup U_2$ for some nonempty open U_1, U_2 , then C must be infinite dimensional.
- 2 X_α possesses Haver's property C (hence, **weakly infinite dimensional**) for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level Borel isomorphic** to X_β .
- 4 If $\alpha \neq \beta$, then the Banach algebra $\mathcal{B}_n^*(X_\alpha)$ is **not linearly isometric** (not ring isomorphic etc.) to $\mathcal{B}_n^*(X_\beta)$ for any $n \in \omega$.

Decomposition Theorem (K.; Gregoriades and K.; K. and Ng)

If $f : X \rightarrow Y$ is a function from analytic sp. X into Polish sp. Y s.t.

$$A \subseteq \Sigma_{\sim m+1}^0(Y) \Rightarrow f^{-1}[A] \in \Sigma_{\sim n+1}^0(X)$$

then, there exists a countable partition $(X_i)_{i \in \omega}$ of X such that the restriction $f|_{X_i}$ is $\Sigma_{\sim n-m+1}^0$ -measurable for every $i \in \omega$.

Decomposition Theorem (K.; Gregoriades and K.; K. and Ng)

If $f : X \rightarrow Y$ is a function from analytic sp. X into Polish sp. Y s.t.

$$A \subseteq \Sigma_{\sim m+1}^0(Y) \Rightarrow f^{-1}[A] \in \Sigma_{\sim n+1}^0(X)$$

then, there exists a countable partition $(X_i)_{i \in \omega}$ of X such that the restriction $f|_{X_i}$ is $\Sigma_{\sim n-m+1}^0$ -measurable for every $i \in \omega$.

Recursion Theoretic Proof

- By the [Louveau separation theorem](#), we have a Borel measurable transition of a $\Sigma_{\sim m+1}^0$ -code of A into a $\Sigma_{\sim n+1}^0$ -code of $f^{-1}[A]$.

Decomposition Theorem (K.; Gregoriades and K.; K. and Ng)

If $f : X \rightarrow Y$ is a function from analytic sp. X into Polish sp. Y s.t.

$$A \subseteq \Sigma_{\sim m+1}^0(Y) \Rightarrow f^{-1}[A] \in \Sigma_{\sim n+1}^0(X)$$

then, there exists a countable partition $(X_i)_{i \in \omega}$ of X such that the restriction $f|_{X_i}$ is $\Sigma_{\sim n-m+1}^0$ -measurable for every $i \in \omega$.

Recursion Theoretic Proof

- By the [Louveau separation theorem](#), we have a Borel measurable transition of a $\Sigma_{\sim m+1}^0$ -code of A into a $\Sigma_{\sim n+1}^0$ -code of $f^{-1}[A]$.
- We then have $(f(x) \oplus z)^{(m)} \leq_T (x \oplus (z \oplus p)^{(\xi)})^{(n)}$ for all $z \in 2^\omega$, where \leq_T is [generalized Turing reducibility](#) on represented spaces.

Decomposition Theorem (K.; Gregoriades and K.; K. and Ng)

If $f : X \rightarrow Y$ is a function from analytic sp. X into Polish sp. Y s.t.

$$A \subseteq \Sigma_{\sim m+1}^0(Y) \Rightarrow f^{-1}[A] \in \Sigma_{\sim n+1}^0(X)$$

then, there exists a countable partition $(X_i)_{i \in \omega}$ of X such that the restriction $f|_{X_i}$ is $\Sigma_{\sim n-m+1}^0$ -measurable for every $i \in \omega$.

Recursion Theoretic Proof

- By the [Louveau separation theorem](#), we have a Borel measurable transition of a $\Sigma_{\sim m+1}^0$ -code of A into a $\Sigma_{\sim n+1}^0$ -code of $f^{-1}[A]$.
- We then have $(f(x) \oplus z)^{(m)} \leq_T (x \oplus (z \oplus p)^{(\xi)})^{(n)}$ for all $z \in 2^\omega$, where \leq_T is [generalized Turing reducibility](#) on represented spaces.
- By the [Shore-Slaman join theorem](#) for any Polish degree structure, we have $f(x) \leq_T (x \oplus p)^{(\xi)(n-m)}$.

Decomposition Theorem (K.; Gregoriades and K.; K. and Ng)

If $f : X \rightarrow Y$ is a function from analytic sp. X into Polish sp. Y s.t.

$$A \subseteq \Sigma_{\sim m+1}^0(Y) \Rightarrow f^{-1}[A] \in \Sigma_{\sim n+1}^0(X)$$

then, there exists a countable partition $(X_i)_{i \in \omega}$ of X such that the restriction $f|_{X_i}$ is $\Sigma_{\sim n-m+1}^0$ -measurable for every $i \in \omega$.

Recursion Theoretic Proof

- By the [Louveau separation theorem](#), we have a Borel measurable transition of a $\Sigma_{\sim m+1}^0$ -code of A into a $\Sigma_{\sim n+1}^0$ -code of $f^{-1}[A]$.
- We then have $(f(x) \oplus z)^{(m)} \leq_T (x \oplus (z \oplus p)^{(\xi)})^{(n)}$ for all $z \in 2^\omega$, where \leq_T is [generalized Turing reducibility](#) on represented spaces.
- By the [Shore-Slaman join theorem](#) for any Polish degree structure, we have $f(x) \leq_T (x \oplus p^{(\xi)})^{(n-m)}$.
- Therefore, f is decomposed into countably many $\Sigma_{\sim n-m+1}^0$ -measurable functions $x \mapsto \Phi_e((x \oplus p^{(\xi)})^{(n-m)})$, $e \in \omega$.

The role of the Decomposition Theorem here is for showing that every n -th Borel isomorphism is covered by ω -many partial homeomorphisms.

$X \leq_{pw} Y$ means that there is a countable cover $\{X_i\}_{i \in \omega}$ of X s.t. X_i is topologically embedded into Y for every $i \in \omega$.

Main Problem

Does there exist an uncountable Polish space X satisfying either of the following equivalent conditions?

- 1 $B_2^*(X)$ is linearly isometric neither to $B_2^*(\mathbb{R})$ nor to $B_2^*(\mathbb{R}^{\mathbb{N}})$.
- 2 $B_2^*(X)$ is ring isomorphic neither to $B_2^*(\mathbb{R})$ nor to $B_2^*(\mathbb{R}^{\mathbb{N}})$.
- 3 X is 2^{nd} level Borel isomorphic neither to \mathbb{R} nor to $\mathbb{R}^{\mathbb{N}}$.
- 4 $\mathbb{R} \prec_{pw} X \prec_{pw} \mathbb{R}^{\mathbb{N}}$.

Compared to the Borel isomorphism problem in 1970s:

- The *Borel isomorphism problem* on analytic spaces was able to be reduced to the same problem on *zero-dimensional* analytic spaces.
- The *second-level Borel isomorphism problem* is inescapably tied to *infinite dimensional* topology.

- Recall: Jayne-Rogers (1979) showed that any two uncountable Polish spaces that are countable unions of sets of finite dimension are 2^{nd} -level Borel isomorphic.
- Indeed, Hurewicz-Wallman (1941) showed that

$$X \simeq_{pw} \mathbb{R} \iff \mathbf{trind}(X) < \infty,$$

where **trind** is transfinite inductive dimension.

- (Alexandrov 1948) X is *weakly infinite dimensional* (w.i.d.) if for each sequence (A_i, B_i) of pairs of disjoint closed sets in X there are separations L_i in X of A_i and B_i s.t. $\bigcap_i L_i = \emptyset$.
- (Haver 1973, Addis-Gresham 1978) X is a **C-space** ($\mathbf{S}_c(O, O)$) if for each sequence (\mathcal{U}_i) of open covers of X there is a pairwise disjoint open family (\mathcal{V}_i) refining (\mathcal{U}_i) s.t. $\bigcup_i \mathcal{V}_i$ covers X .

$X \leq_{pw} 2^{\mathbb{N}} \Leftrightarrow \text{trind}(X) < \infty \Rightarrow X \text{ is } \mathbf{C} \Rightarrow X \text{ is w.i.d.}$

- (Alexandrov 1951) \exists a w.i.d. metrizable compactum $X \succ_{pw} 2^{\mathbb{N}}?$
- (R. Pol 1981) There exists a metrizable **C**-compactum $X \succ_{pw} 2^{\mathbb{N}}$.
- (E. Pol 1997) There exists an infinite dimensional **C**-Cantor manifold, i.e., a **C**-compactum which cannot be separated by any hereditarily weakly infinite dimensional closed subspaces.
- (Chatyrko 1999) There is a collection $\{X_\alpha\}_{\alpha < 2^{\aleph_0}}$ of continuum many infinite dimensional **C**-Cantor manifolds such that X_α cannot be embedded into X_β whenever $\alpha \neq \beta$.

An *infinite dimensional \mathbf{C} -Cantor manifold* is a \mathbf{C} -compactum which cannot be separated by any hereditarily weakly infinite dimensional closed subspace.


Main Lemma (K. and Pauly)

Let \mathfrak{M}_∞ be the class of all infinite dimensional \mathbf{C} -Cantor manifolds. Then, there is an order embedding of $([\mathfrak{N}_1]^\omega, \subseteq)$ into $(\mathfrak{M}_\infty, \leq_{pw})$.


- This solves Motto Ros' problem (and the second level Borel isomorphism problem) in Banach Space Theory.
- This strengthens R. Pol's theorem and Chatyrko's theorem in Infinite Dimensional Topology.

To show Main Lemma, we again use **Recursion Theory!**

Idea of Proof: Upper/Lower Approximation by Zero Dim Spaces




(a) Any point in \mathbb{R}^n




(b) Some point in $[0, 1]^{\mathbb{N}}$

Idea of Proof: Upper/Lower Approximation by Zero Dim Spaces




(a) Any point in \mathbb{R}^n



(b) Some point in $[0, 1]^{\mathbb{N}}$

- By *approximating* each point in a space X by a zero-dim space, we measure “*how similar the space X is to a zero-dim space*”.
- (a) Upper and lower approximations by a zero-dim space *meet*.
- (b) There is a *gap* between upper and lower approximations by a zero-dim space

Idea of Proof: Upper/Lower Approximation by Zero Dim Spaces



- $\text{Spec}(x) = \{p \in 2^{\mathbb{N}} : x \leq_T p\}$.
- $\text{coSpec}(x) = \{p \in 2^{\mathbb{N}} : p \leq_T x\}$.

Key Idea

Classification of topological spaces by degrees of unsolvability:

- 1 The Turing degrees \simeq the degree structure on Cantor space $2^{\mathbb{N}}$ and Euclidean spaces \mathbb{R}^n .
- 2 The enumeration degrees \simeq the degree structure on the Scott domain $\mathcal{P}(\mathbb{N})$.
- 3 Hinman (1973): degrees of unsolvability of continuous functionals \simeq the degree structure on the space $\mathbb{N}^{\mathbb{N}^{\mathbb{N}}}$ of Kleene-Kreisel continuous functionals.
- 4 J. Miller (2004): continuous degrees \simeq the degree structure on the function space $C([0, 1])$ and the Hilbert cube $[0, 1]^{\mathbb{N}}$.

Definition

Let X and Y be second-countable T_0 spaces with fixed countable open basis $\{B_n^X\}_{n \in \omega}$ and $\{B_n^Y\}_{n \in \omega}$.

A point $x \in X$ is “*Turing reducible*” to a point $y \in Y$ ($x \leq_T y$) if


$$\{n \in \omega : x \in B_n^X\} \leq_e \{n \in \omega : y \in B_n^Y\}.$$

In other words, we identify the “*Turing degree*” of $x \in X$ with the *enumeration degree of the (coded) neighborhood filter* of x .

Example

- The degree structure of *Cantor space* is exactly the same as the *Turing degrees*.
- The degree structure of *Hilbert cube* (a universal Polish space) is exactly the same as the *continuous degrees*.
- The degree structure of *the Scott domain $O(\mathbb{N})$* (a universal quasi-Polish space) is exactly the same as the *enumeration degrees*.

Idea of Proof: Upper/Lower Approximation by Zero Dim Spaces



(a) Any point in \mathbb{R}^n

(b) Some point in $[0, 1]^{\mathbb{N}}$

- $\text{Spec}(x) = \{p \in 2^{\mathbb{N}} : x \leq_T p\}$.
- $\text{coSpec}(x) = \{p \in 2^{\mathbb{N}} : p \leq_T x\}$.

$\text{Spec}(x) = \{\rho \in 2^{\mathbb{N}} : x \leq_T \rho\}; \text{Spec}(X) = \{\text{Spec}(x) : x \in X\}.$
 $\text{coSpec}(x) = \{\rho \in 2^{\mathbb{N}} : \rho \leq_T x\}; \text{coSpec}(X) = \{\text{coSpec}(x) : x \in X\}$

$\text{Spec}(x) = \{p \in 2^{\mathbb{N}} : x \leq_T p\}$; $\text{Spec}(X) = \{\text{Spec}(x) : x \in X\}$.
 $\text{coSpec}(x) = \{p \in 2^{\mathbb{N}} : p \leq_T x\}$; $\text{coSpec}(X) = \{\text{coSpec}(x) : x \in X\}$

Lemma (K. and Pauly)

$X \simeq_{pw} Y \implies \text{Spec}^r(X) = \text{Spec}^r(Y)$ for some oracle $r \in 2^\omega$.
 $\implies \text{coSpec}^r(X) = \text{coSpec}^r(Y)$ for some oracle $r \in 2^\omega$.

$\text{Spec}(x) = \{p \in 2^{\mathbb{N}} : x \leq_T p\}$; $\text{Spec}(X) = \{\text{Spec}(x) : x \in X\}$.
 $\text{coSpec}(x) = \{p \in 2^{\mathbb{N}} : p \leq_T x\}$; $\text{coSpec}(X) = \{\text{coSpec}(x) : x \in X\}$

Lemma (K. and Pauly)


$X \simeq_{pw} Y \implies \text{Spec}^r(X) = \text{Spec}^r(Y)$ for some oracle $r \in 2^\omega$.
 $\implies \text{coSpec}^r(X) = \text{coSpec}^r(Y)$ for some oracle $r \in 2^\omega$.

- 1 A Turing ideal $\mathcal{J} \subseteq 2^\omega$ is *realized* by x if $\mathcal{J} = \text{coSpec}(x)$.
- 2 A countable set $\mathcal{J} \subseteq 2^\omega$ is a *Scott ideal*
 $\iff (\omega, \mathcal{J}) \models \text{RCA} + \text{WKL}$.

Realizability of Scott ideals (J. Miller 2004)

- 1 $2^\omega \simeq_{pw} \omega^\omega \simeq_{pw} \mathbb{R}^n \simeq_{pw} \bigoplus_{n \in \omega} \mathbb{R}^n$. (*Turing degrees.*)
No Scott ideal is realized in these spaces!
- 2 $[0, 1]^\omega \simeq_{pw} C([0, 1]) \simeq_{pw} \ell^2$. (*full continuous degrees.*)
Every countable Scott ideal is realized in these spaces!

Idea of Proof: Upper/Lower Approximation by Zero Dim Spaces



(a) Any point in \mathbb{R}^n

(b) Some point in $[0, 1]^{\mathbb{N}}$

- **Spec** determines the pw-homeomorphism type of a space, and **coSpec** is invariant under pw-homeomorphism.
- The **coSpec** of any point in a space of **dim** $< \infty$ has to be a principal Turing ideal.
- (Miller) Every countable Scott ideal is realized as **coSpec** of a point in Hilbert cube.

Definition

$\Gamma : 2^{\mathbb{N}} \rightarrow [0, 1]^{\mathbb{N}}$ is *ω -left-CEA operator* if the infinite sequence $\Gamma(\mathbf{y}) = (\mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots)$ is generated in a uniformly left-computably enumerable manner by a single Turing machine, that is, there is a left-c.e. operator γ such that for all i ,

$$\mathbf{x}_i := \Gamma(\mathbf{y})(i) = \gamma(\mathbf{y}, i, \mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{i-1}).$$

An ω -left-CEA operator $\Gamma : \mathbb{N} \times 2^{\mathbb{N}} \rightarrow [0, 1]^{\mathbb{N}}$ is *universal* if for every ω -left-CEA operator Ψ , there is \mathbf{e} such that $\Psi = \lambda \mathbf{y}. \Gamma(\mathbf{e}, \mathbf{y})$.

Let ωCEA denote the graph of a universal ω -left-CEA operator.

Theorem (K.-Pauly)


The space ωCEA (as a subspace of Hilbert cube) is an intermediate Polish space:

$$2^{\mathbb{N}} \prec_{pw} \omega\text{CEA} \prec_{pw} [0, 1]^{\mathbb{N}}$$

Remark

Furthermore, ωCEA is pw-homeomorphic to the following:

- Rubin-Schori-Walsh (1979)'s **strongly infinite dimensional totally disconnected Polish space**.
- Roman Pol (1981)'s **weakly infinite dimensional compactum which is not decomposable into countably many finite-dim subspaces** (a solution to Alexandrov's problem).



(a) $2^{\mathbb{N}}$

(b) ωCEA

(c) $[0, 1]^{\mathbb{N}}$

- (a) **coSpec** is principal, and *meets* with **Spec**.
- (b) **coSpec** is not always principal, but the “*distance*” between **Spec** and **coSpec** has to be at most *the ω -th Turing jump*.
- (c) **coSpec** can realize an arbitrary countable Scott ideal, hence **Spec** and **coSpec** can be separated by *an arbitrary distance*.

$\omega\mathbf{CEA} = \{(\mathbf{e}, \mathbf{p}, \mathbf{x}_0, \mathbf{x}_1, \dots) \in \omega \times 2^\omega \times [0, 1]^\omega :$
 $(\forall i) \mathbf{x}_i \text{ is the } \mathbf{e}\text{-th left-c.e. real in } (\mathbf{p}, \mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{i-1}).\}$

Lemma

For any $\mathbf{p} \in 2^\omega$, the following Scott ideal is not realized in $\omega\mathbf{CEA}$:

$$\mathcal{J}^{\mathbf{p}} = \{\mathbf{z} \in 2^\omega : (\exists n) \mathbf{z} \leq_T \mathbf{p}^{(\omega \cdot n)}\}.$$

- Pick $\mathbf{z} = (\mathbf{e}, \mathbf{p}, \mathbf{x}_0, \mathbf{x}_1, \dots) \in \omega\mathbf{CEA}$.
- Then, $\mathbf{p} \in \mathbf{coSpec}(\mathbf{z})$ and $\mathbf{p}^{(\omega)} \in \mathbf{Spec}(\mathbf{z})$.
- Clearly, $\mathbf{p}^{(\omega+1)} \notin \mathbf{coSpec}(\mathbf{z})$.

Since \mathbf{coSpec} (up to an oracle) is invariant under pw-homeomorphism, we have $\omega\mathbf{CEA} \prec_{pw} [0, 1]^{\mathbb{N}}$.

Another separation is based on Kakutani's fixed point theorem.


Theorem (J. Miller 2004)

There is a nonempty convex-valued computable function $\Psi : [0, 1]^{\mathbb{N}} \rightarrow \mathcal{P}([0, 1]^{\mathbb{N}})$ with a closed graph such that for every fixed point $\langle \mathbf{x}_0, \mathbf{x}_1, \dots \rangle \in \mathbf{Fix}(\Psi)$,

$$\mathbf{coSpec}(\langle \mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots \rangle) = \{\mathbf{x}_0, \mathbf{x}_1, \mathbf{x}_2, \dots\}.$$

Moreover, such an \mathbf{x} realizes a Scott ideal.

- $\mathbf{Fix}(\Psi)$ is a Π_1^0 subset of $[0, 1]^{\omega}$.
- Inductively find $(\mathbf{x}_0, \mathbf{x}_1, \dots) \in \mathbf{Fix}(\Psi)$, where \mathbf{x}_{i+1} is the “leftmost” value s.t. $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_{i+1})$ is extendible in $\mathbf{Fix}(\Psi)$.
- Then, \mathbf{x}_{i+1} is left-c.e. in $(\mathbf{x}_0, \mathbf{x}_1, \dots, \mathbf{x}_i)$, uniformly.
- \mathbf{x}_{i+1} does not depend on the choice of a name of $(\mathbf{x}_0, \dots, \mathbf{x}_i)$.



(a) $2^{\mathbb{N}}$

(b) ωCEA

(c) $[0, 1]^{\mathbb{N}}$

- (a) **coSpec** is principal, and *meets* with **Spec**.
- (b) **coSpec** is not always principal, but the “*distance*” between **Spec** and **coSpec** has to be at most *the ω -th Turing jump*.
- (c) **coSpec** can realize an arbitrary countable Scott ideal, hence **Spec** and **coSpec** can be separated by *an arbitrary distance*.

- 1 $\mathbf{coSpec}(2^{\mathbb{N}})$ = all principal Turing ideals.
- 2 $\mathbf{coSpec}([0, 1]^{\mathbb{N}})$ = all principal Turing ideals and Scott ideals.
- 3 What do we know about $\mathbf{coSpec}(\omega\mathbf{CEA})$?
 - It cannot realize an ω -jump ideal.
 - It realizes a non-principal Turing ideal.
 - We know absolutely nothing about what kind of Turing ideals it realizes; even whether it realizes a jump ideal or not.

How can we control \mathbf{coSpec} of a Polish space?

For instance, given $\alpha \ll \beta < \omega_1$, we need a technique for constructing a Polish space such that

- it cannot realize a β -jump ideal,
- it realizes an α -jump ideal.

We say that $\mathcal{G} : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ is an *oracle Π_2^0 singleton* if it has a Π_2^0 graph. For instance, the α -th Turing jump operator \mathbf{TJ}^α is an oracle Π_2^0 singleton.

Definition (Modified ω CEA Space)

The space $\omega\mathbf{CEA}(\mathcal{G})$ consists of $(\mathbf{d}, \mathbf{e}, \mathbf{r}, \mathbf{x}) \in \mathbb{N}^2 \times 2^{\mathbb{N}} \times [0, 1]^{\mathbb{N}}$ such that for every i ,

- ① either $x_i = \mathcal{G}^i(\mathbf{r})$, or
- ② there are $u \leq v \leq i$ such that $x_i \in [0, 1]$ is the \mathbf{e} -th left-c.e. real in $\langle \mathbf{r}, \mathbf{x}_{<i}, \mathbf{x}_{I(u)} \rangle$ and $x_{I(u)} = \mathcal{G}^{I(u)}(\mathbf{r})$, where $I(u) = \Phi_{\mathbf{d}}(u, \mathbf{r}, \mathbf{x}_{<v})$.

Here: $\mathcal{G}^0(\mathbf{x}) = \mathbf{x}$ and $\mathcal{G}^{n+1}(\mathbf{x}) = \mathcal{G}^n(\mathbf{x}) \oplus \mathcal{G}(\mathcal{G}^n(\mathbf{x}))$.

We define $\mathbf{Ref}(\mathcal{G}) = \omega\mathbf{CEA}(\mathcal{G}) \cap (\mathbb{N}^2 \times \mathbf{Fix}(\Psi))$.

The subspace $\mathbf{Ref}(\mathcal{G})$ (as a subspace of $[0, 1]^{\mathbb{N}}$) is Polish whenever \mathcal{G} is an oracle Π_2^0 singleton.

Suppose that \mathcal{G} is an oracle Π_2^0 -singleton. For every oracle $r \in 2^{\mathbb{N}}$, consider two Turing ideals defined as

$$\begin{aligned} \mathcal{J}_T(\mathcal{G}, r) &= \{z \in 2^{\mathbb{N}} : (\exists n \in \mathbb{N}) x \leq_T \mathcal{G}^n(r)\}, \\ \mathcal{J}_a(\mathcal{G}, r) &= \{z \in 2^{\mathbb{N}} : (\exists n \in \mathbb{N}) x \leq_a \mathcal{G}^n(r)\}. \end{aligned}$$

Here: \leq_a is the arithmetical reducibility.

Main Lemma (coSpec-Controlling)

- 1 For every $x \in \mathbf{Ref}(\mathcal{G})$, there is $r \in 2^{\mathbb{N}}$ such that

$$\mathbf{coSpec}(x) \subseteq \mathcal{J}_a(\mathcal{G}, r).$$
- 2 For every $r \in 2^{\mathbb{N}}$, there is $x \in \mathbf{Ref}(\mathcal{G})$ such that

$$\mathcal{J}_T(\mathcal{G}, r) \subseteq \mathbf{coSpec}(x).$$

If $\mathcal{G} = \mathbf{TJ}^\alpha$ is the α -th Turing jump operator for $\alpha \geq \omega$,

- 1 $\mathbf{coSpec}(\mathbf{Ref}(\mathbf{TJ}^\alpha))$ realizes no β -jump ideal for $\beta \geq \alpha \cdot \omega$,
- 2 $\mathbf{coSpec}(\mathbf{Ref}(\mathbf{TJ}^\alpha))$ realizes an α -jump ideal.

- 1 By **coSpec**-Controlling Lemma, given an oracle Π_2^0 singleton \mathcal{G} we can construct a Polish space which realizes all Turing ideals closed under \mathcal{G} .
- 2 $\mathbf{Ref}(\mathcal{G})$ is strongly infinite dimensional and totally disconnected.
- 3 Hence, its compactification $\gamma\mathbf{Ref}(\mathcal{G})$ (in the sense of Lelek) is a “*Pol-type space*”, hence, a metrizable \mathbf{C} -compacta.
- 4 Note that Lelek’s compactification preserves **Spec** and **coSpec**.
- 5 By combining Elzbieta Pol’s construction, our spaces can be assumed to be infinite dimensional \mathbf{C} -Cantor manifolds.

Main Lemma (K. and Pauly)

Let \mathfrak{M}_∞ be the class of all infinite dimensional \mathbf{C} -Cantor manifolds. Then, there is an order embedding of $([\mathbb{N}_1]^\omega, \subseteq)$ into $(\mathfrak{M}_\infty, \leq_{pw})$.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$,
- 2 X_α possesses Haver's property **C** for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level isomorphic** to X_β for any $n \in \omega$.
- 4 If $\alpha \neq \beta$, then the Banach space $\mathcal{B}_n(X_\alpha)$ is **not linearly isometric** to $\mathcal{B}_n(X_\beta)$ for any $n \in \omega$.

Summary of This Work

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$,
- 2 X_α possesses Haver's property **C** for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level isomorphic** to X_β for any $n \in \omega$.
- 4 If $\alpha \neq \beta$, then the Banach space $\mathcal{B}_n(X_\alpha)$ is **not linearly isometric** to $\mathcal{B}_n(X_\beta)$ for any $n \in \omega$.

Summary of This Work

- 1 Defining the notion of **Spec** and **coSpec**.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$,
- 2 X_α possesses Haver's property **C** for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level isomorphic** to X_β for any $n \in \omega$.
- 4 If $\alpha \neq \beta$, then the Banach space $\mathcal{B}_n(X_\alpha)$ is **not linearly isometric** to $\mathcal{B}_n(X_\beta)$ for any $n \in \omega$.

Summary of This Work

- 1 Defining the notion of **Spec** and **coSpec**.
- 2 Using **Spec** and **coSpec** as "*pw-topological*" invariant.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$,
- 2 X_α possesses Haver's property **C** for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level isomorphic** to X_β for any $n \in \omega$.
- 4 If $\alpha \neq \beta$, then the Banach space $\mathcal{B}_n(X_\alpha)$ is **not linearly isometric** to $\mathcal{B}_n(X_\beta)$ for any $n \in \omega$.

Summary of This Work

- 1 Defining the notion of **Spec** and **coSpec**.
- 2 Using **Spec** and **coSpec** as "*pw-topological*" invariant.
- 3 Proving **coSpec**-Controlling Lemma.

Main Theorem (K. and Pauly)

There exists a 2^{\aleph_0} collection $(X_\alpha)_{\alpha < 2^{\aleph_0}}$ of topological spaces s.t.

- 1 X_α is an infinite dimensional Cantor manifold for any $\alpha < 2^{\aleph_0}$,
- 2 X_α possesses Haver's property **C** for any $\alpha < 2^{\aleph_0}$.
- 3 If $\alpha \neq \beta$, then X_α is **not n -th level isomorphic** to X_β for any $n \in \omega$.
- 4 If $\alpha \neq \beta$, then the Banach space $\mathcal{B}_n(X_\alpha)$ is **not linearly isometric** to $\mathcal{B}_n(X_\beta)$ for any $n \in \omega$.

Summary of This Work

- 1 Defining the notion of **Spec** and **coSpec**.
- 2 Using **Spec** and **coSpec** as "*pw-topological*" invariant.
- 3 Proving **coSpec**-Controlling Lemma.
- 4 Solving the second-level Borel isomorphism problem.

Open Question

- 1 What is the role of $\mathcal{B}_2(\mathbf{X})$ in Banach Space Theory?
 - Note: $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} has a great role in Banach Space Theory, in particular, in the context of *Rosenthal's ℓ^1 Theorem*. A compact subspace of $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} is known as a *Rosenthal compactum*.

Open Question

- 1 What is the role of $\mathcal{B}_2(\mathbf{X})$ in Banach Space Theory?
 - Note: $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} has a great role in Banach Space Theory, in particular, in the context of *Rosenthal's ℓ^1 Theorem*. A compact subspace of $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} is known as a *Rosenthal compactum*.
- 2 Does there exist a *strongly* infinite dimensional compactum \mathbf{X} s.t. $\mathbb{R} \prec_{pw} \mathbf{X} \prec_{pw} \mathbb{R}^{\mathbb{N}}$?
 - Our spaces are all *Pol-type spaces*.

Open Question

- 1 What is the role of $\mathcal{B}_2(\mathbf{X})$ in Banach Space Theory?
 - Note: $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} has a great role in Banach Space Theory, in particular, in the context of *Rosenthal's ℓ^1 Theorem*. A compact subspace of $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} is known as a *Rosenthal compactum*.
- 2 Does there exist a *strongly* infinite dimensional compactum \mathbf{X} s.t. $\mathbb{R} \prec_{pw} \mathbf{X} \prec_{pw} \mathbb{R}^{\mathbb{N}}$?
 - Our spaces are all *Pol-type spaces*.
- 3 Develop the degree theory on non-second-countable spaces.
 - e.g., the space $\mathbb{N}^{\mathbb{N}^{\mathbb{N}}}$ of Kleene-Kreisel continuous functionals.
 - computability theory on the spaces $\mathcal{B}_\alpha(\mathbf{X})$?

Open Question

- 1 What is the role of $\mathcal{B}_2(\mathbf{X})$ in Banach Space Theory?
 - Note: $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} has a great role in Banach Space Theory, in particular, in the context of *Rosenthal's ℓ^1 Theorem*. A compact subspace of $\mathcal{B}_1(\mathbf{X})$ for Polish \mathbf{X} is known as a *Rosenthal compactum*.
- 2 Does there exist a *strongly* infinite dimensional compactum \mathbf{X} s.t. $\mathbb{R} \prec_{pw} \mathbf{X} \prec_{pw} \mathbb{R}^{\mathbb{N}}$?
 - Our spaces are all *Pol-type spaces*.
- 3 Develop the degree theory on non-second-countable spaces.
 - e.g., the space $\mathbb{N}^{\mathbb{N}^{\mathbb{N}}}$ of Kleene-Kreisel continuous functionals.
 - computability theory on the spaces $\mathcal{B}_\alpha(\mathbf{X})$?
- 4 Develop the notion of a hyperdegree spectrum of a space.
 - Gregoriades and K. have already studied co-Souslin- \mathcal{F} isomorphisms as counterpart of hyperdegree spectra, and obtained a few results based on classical works on the Borel isomorphism problem, Kleene degrees (real computability relative to ${}^2\mathbf{E}$) and so on.