

Parametric Approximation of Geometric Evolution Equations

John W. Barrett,

Department of Mathematics, Imperial College, London, SW7 2AZ, UK

joint work with

Harald Garcke (Universität Regensburg, Germany)

and

Robert Nürnberg (Imperial College)

Geometric flows, in which hypersurfaces move such that an energy, involving surface and bending terms, decreases appear in many situations in the natural sciences and in geometry. Classic examples are mean curvature, surface diffusion and Willmore flows. Computational methods to approximate such flows are based on one of three approaches (i) parametric methods, (ii) phase field methods or (iii) level set methods. The first tracks the hypersurface, whilst the other two implicitly capture the hypersurface. A key problem with the first approach, apart from the fact that it does not naturally deal with changes of topology, is that in many cases the mesh has to be redistributed after every few time steps to avoid coalescence of mesh points.

In this talk we present a new variational formulation of the parametric approach, which leads to an unconditionally stable, fully discrete approximation. In addition, the scheme has very good properties with respect to the distribution of mesh points, and if applicable volume conservation. We illustrate this for (anisotropic) mean curvature, (anisotropic) surface diffusion, and Willmore flows of closed curves in \mathbb{R}^2 . We extend the first two of these flows to curve networks in \mathbb{R}^2 . Here the triple junction conditions, that have to hold where three curves meet at a point, are naturally approximated in the discretization of our variational formulation. Finally, we extend these flows to closed hypersurfaces in \mathbb{R}^3 .