

Weak Degrees of Π_1^0 Subsets of 2^ω

Stephen G. Simpson

Pennsylvania State University
<http://www.math.psu.edu/simpson/>
simpson@math.psu.edu

Computational Prospects of Infinity
Institute for Mathematical Sciences
National University of Singapore
June 20 – August 15, 2005

Motivation:

Recall that \mathcal{R}_T is the upper semilattice of recursively enumerable Turing degrees.

Two basic, classical, unresolved issues concerning \mathcal{R}_T are:

Issue 1. To find a specific, natural, r.e. Turing degree $\mathbf{a} \in \mathcal{R}_T$ which is $> \mathbf{0}$ and $< \mathbf{0}'$.

Issue 2. To find a “smallness property” of an infinite co-r.e. set $A \subseteq \omega$ which ensures that $\deg_T(A) = \mathbf{a} \in \mathcal{R}_T$ is $> \mathbf{0}$ and $< \mathbf{0}'$.

These unresolved issues go back to Post's 1944 paper, *Recursively enumerable sets of positive integers and their decision problems*.

Mass Problems to the Rescue!

We address Issues 1 and 2 by passing from decision problems to mass problems.

Outline of this talk:

We embed \mathcal{R}_T into a slightly larger structure, \mathcal{P}_w , which is much better behaved. In the \mathcal{P}_w context, we obtain satisfactory, positive answers to Issues 1 and 2.

What is this wonderful structure \mathcal{P}_w ?

Briefly, \mathcal{P}_w is the lattice of weak degrees of mass problems associated with nonempty Π_1^0 subsets of 2^ω .

In order to explain \mathcal{P}_w , we must first explain:

- mass problems,
- weak degrees, and
- nonempty Π_1^0 subsets of 2^ω .

Mass problems (informal discussion):

A “decision problem” is the problem of deciding whether a given $n \in \omega$ belongs to a fixed set $A \subseteq \omega$ or not. To compare decision problems, we use Turing reducibility. $A \leq_T B$ means that A can be computed using an oracle for B .

A “mass problem” is a problem with a not necessarily unique solution. (By contrast, a “decision problem” has only one solution.)

The “mass problem” associated with a set $P \subseteq \omega^\omega$ is the “problem” of computing an element of P .

The “solutions” of P are the elements of P .

One mass problem is said to be “reducible” to another if, given any solution of the second problem, we can use it as an oracle to compute a solution of the first problem.

Rigorous definition:

Let P and Q be subsets of ω^ω .

We view P and Q as mass problems.

We say that P is *weakly reducible* to Q if

$$(\forall Y \in Q) (\exists X \in P) (X \leq_T Y) .$$

This is abbreviated $P \leq_w Q$.

Summary:

$P \leq_w Q$ means that, given any solution of Q , we can use it as an oracle to compute a solution of P .

Digression: weak vs. strong reducibility

Let P and Q be subsets of ω^ω .

1. P is *weakly reducible* to Q , $P \leq_w Q$, if for all $Y \in Q$ there exists e such that $\{e\}^Y \in P$.
2. P is *strongly reducible* to Q , $P \leq_s Q$, if there exists e such that $\{e\}^Y \in P$ for all $Y \in Q$.

Strong reducibility is a uniform variant of weak reducibility. By a result of Nerode, there is an analogy:

$$\frac{\text{weak reducibility}}{\text{Turing reducibility}} = \frac{\text{strong reducibility}}{\text{truth table reducibility}} .$$

In this talk we deal only with weak reducibility.

Historical note:

Weak reducibility is due to Muchnik 1963.

Strong reducibility is due to Medvedev 1955.

The lattice \mathcal{P}_w :


We focus on Π_1^0 subsets of 2^ω , i.e.,
 $P = \{\text{paths through } T\}$ where T is a recursive subtree of $2^{<\omega}$, the full binary tree of finite sequences of 0's and 1's.

We define \mathcal{P}_w to be the set of weak degrees of nonempty Π_1^0 subsets of 2^ω , ordered by weak reducibility.

Basic facts about \mathcal{P}_w :

1. \mathcal{P}_w is a distributive lattice, with l.u.b. given by $P \times Q = \{X \oplus Y \mid X \in P, Y \in Q\}$, and g.l.b. given by $P \cup Q$.
2. The bottom element of \mathcal{P}_w is the weak degree of 2^ω .
3. The top element of \mathcal{P}_w is the weak degree of $\text{PA} = \{\text{completions of Peano Arithmetic}\}$. (Scott/Tennenbaum).

Weak reducibility of Π_1^0 subsets of 2^ω :



$P \leq_w Q$ means:

$$(\forall Y \in Q) (\exists X \in P) (X \leq_T Y).$$

P, Q are given by infinite recursive subtrees of the full binary tree of finite sequences of 0's and 1's.

X, Y are infinite (nonrecursive) paths through P, Q respectively.

The lattice \mathcal{P}_w (review):

A *weak degree* is an equivalence class of subsets of ω^ω under the equivalence relation $P \leq_w Q$ and $Q \leq_w P$. The weak degrees have a partial ordering induced by \leq_w .

We define \mathcal{P}_w to be the set of weak degrees of nonempty Π_1^0 subsets of 2^ω , partially ordered by weak reducibility.

\mathcal{P}_w is a countable distributive lattice.

The bottom element of \mathcal{P}_w is the weak degree of 2^ω .

The top element of \mathcal{P}_w is the weak degree of $\text{PA} = \{\text{completions of Peano Arithmetic}\}$.

Embedding \mathcal{R}_T into \mathcal{P}_w :

Theorem (Simpson 2002):

There is a natural embedding $\phi : \mathcal{R}_T \rightarrow \mathcal{P}_w$.

($\mathcal{R}_T =$ the semilattice of Turing degrees of r.e. subsets of ω . $\mathcal{P}_w =$ the lattice of weak degrees of nonempty Π_1^0 subsets of 2^ω .)

The embedding ϕ is given by

$$\phi : \text{deg}_T(A) \mapsto \text{deg}_w(\text{PA} \cup \{A\}).$$

Note: $\text{PA} \cup \{A\}$ is not a Π_1^0 set. However, it is of the same weak degree as a Π_1^0 set. This is already a nontrivial result. See below.


The embedding ϕ is one-to-one and preserves \leq , l.u.b., and the top and bottom elements.

Convention:

We identify \mathcal{R}_T with its image in \mathcal{P}_w under ϕ .

In particular, we identify $\mathbf{0}', \mathbf{0} \in \mathcal{R}_T$ with the top and bottom elements of \mathcal{P}_w .

A picture of the lattice \mathcal{P}_w :



\mathcal{R}_T is embedded in \mathcal{P}_w . $0'$ and 0 are the top and bottom elements of both \mathcal{R}_T and \mathcal{P}_w .

Structural properties of \mathcal{P}_w :

1. \mathcal{P}_w is a countable distributive lattice.
Every countable distributive lattice is lattice embeddable in every initial segment of \mathcal{P}_w .
(Binns/Simpson 2001)

2. The \mathcal{P}_w analog of the Sacks Splitting Theorem holds. (Stephen Binns, 2002)

3. We conjecture that the \mathcal{P}_w analog of the Sacks Density Theorem holds.

These structural results for \mathcal{P}_w are proved by means of priority arguments, just as for \mathcal{R}_T .

4. Within \mathcal{P}_w the degrees \mathbf{r}_1 and $\inf(\mathbf{r}_2, \mathbf{0}')$ are meet irreducible and do not join to $\mathbf{0}'$.
(Simpson 2002, 2004)

5. $\mathbf{0}$ is meet irreducible. (This is trivial.)

Response to Issue 1:

Issue 1 was:


To find a specific, natural example of a recursively enumerable Turing degree which is $> \mathbf{0}$ and $< \mathbf{0}'$.

We do not know how to do this.

However, in the \mathcal{P}_w context, we have discovered many specific, natural degrees which are $> \mathbf{0}$ and $< \mathbf{0}'$.

The specific, natural degrees in \mathcal{P}_w which we have discovered are related to foundationally interesting topics:

- algorithmic randomness,
- diagonal nonrecursiveness,
- reverse mathematics,
- subrecursive hierarchies,
- computational complexity.



Note: Except for $0'$ and 0 , the r.e. Turing degrees are incomparable with all of the known, specific, natural degrees in \mathcal{P}_w .

Some specific, natural degrees in \mathcal{P}_w :

\mathbf{r}_n = the weak degree of the set of n -random reals.

\mathbf{d} = the weak degree of the set of diagonally nonrecursive functions.

\mathbf{d}_{REC} = the weak degree of the set of diagonally nonrecursive functions which are recursively bounded.

Theorem (Simpson 2002, Ambos-Spies et al 2004):

In \mathcal{P}_w we have

$$0 < \mathbf{d} < \mathbf{d}_{\text{REC}} < \mathbf{r}_1 < \inf(\mathbf{r}_2, \mathbf{0}') < \mathbf{0}'.$$

Theorem (Simpson 2004):

1. \mathbf{r}_1 is the maximum weak degree of a Π_1^0 subset of 2^ω which is of positive measure.
2. $\inf(\mathbf{r}_2, \mathbf{0}')$ is the maximum weak degree of a Π_1^0 subset of 2^ω whose Turing upward closure is of positive measure.

Structural properties of \mathcal{P}_w :

1. \mathcal{P}_w is a countable distributive lattice.
Every countable distributive lattice is lattice embeddable in every initial segment of \mathcal{P}_w .
(Binns/Simpson 2001)

2. The \mathcal{P}_w analog of the Sacks Splitting Theorem holds. (Stephen Binns, 2002)

3. We conjecture that the \mathcal{P}_w analog of the Sacks Density Theorem holds.

These structural results for \mathcal{P}_w are proved by means of priority arguments, just as for \mathcal{R}_T .

4. Within \mathcal{P}_w the degrees \mathbf{r}_1 and $\inf(\mathbf{r}_2, \mathbf{0}')$ are meet irreducible and do not join to $\mathbf{0}'$.
(Simpson 2002, 2004)

5. $\mathbf{0}$ is meet irreducible. (This is trivial.)

The Embedding Lemma:

If $S \subseteq \omega^\omega$ is Σ_3^0 and if $P \subseteq 2^\omega$ is Π_1^0 , then $\deg_w(S \cup P) \in \mathcal{P}_w$.

It follows that, for many Σ_3^0 sets $S \subseteq \omega^\omega$, $\deg_w(S) \in \mathcal{P}_w$.

Examples:

1. $R_1 = \{X \in 2^\omega \mid X \text{ is 1-random}\}$.

Since R_1 is Σ_2^0 , it follows by the Embedding Lemma that $\mathbf{r}_1 = \deg_w(R_1) \in \mathcal{P}_w$.

2. $R_2 = \{X \in 2^\omega \mid X \text{ is 2-random}\}$.

Since R_2 is Σ_3^0 , it follows by the Embedding Lemma that $\inf(\mathbf{r}_2, \mathbf{0}') = \deg_w(R_2 \cup PA) \in \mathcal{P}_w$.

3. $D = \{f \in \omega^\omega \mid f \text{ is diagonally nonrecursive}\}$.

Since D is Π_1^0 , $\mathbf{d} = \deg_w(D) \in \mathcal{P}_w$.

4. $D_{\text{REC}} = \{f \in D \mid f \text{ is recursively bounded}\}$.

Since D_{REC} is Σ_3^0 , $\mathbf{d}_{\text{REC}} = \deg_w(D_{\text{REC}}) \in \mathcal{P}_w$.

5. Let $A \subseteq \omega$ be r.e. Since $\{A\}$ is Π_2^0 , $\deg_w(\{A\} \cup PA) \in \mathcal{P}_w$. This gives our embedding of \mathcal{R}_T into \mathcal{P}_w .

The Embedding Lemma (restated):

Let $S \subseteq \omega^\omega$ be Σ_3^0 . Let $P \subseteq 2^\omega$ be nonempty Π_1^0 . Then \exists nonempty Π_1^0 $Q \subseteq 2^\omega$ such that $Q \equiv_w S \cup P$.

Proof (sketch). **Step 1.** By Skolem functions, we may assume that $S \subseteq \omega^\omega$ is Π_1^0 .

Step 2. We have $S = \{\text{paths through } T_S\}$, $P = \{\text{paths through } T_P\}$, where T_S, T_P are recursive subtrees of $\omega^{<\omega}, 2^{<\omega}$ respectively. May assume $\tau(n) \geq 2$ for all $n < |\tau|$, $\tau \in T_S$. Define $Q = \{\text{paths through } T_Q\}$, where T_Q is the set of all $\rho \in \omega^{<\omega}$ of the form

$$\rho = \sigma_0 \hat{\ } \langle m_0 \rangle \hat{\ } \sigma_1 \hat{\ } \langle m_1 \rangle \hat{\ } \cdots \hat{\ } \langle m_{k-1} \rangle \hat{\ } \sigma_k$$

where

- $\sigma_0, \sigma_1, \dots, \sigma_k \in T_P$,
- $\langle m_0, m_1, \dots, m_{k-1} \rangle \in T_S$,
- $\rho(n) \leq \max(n, 2)$ for all $n < |\rho|$.

One can show that $Q \equiv_w S \cup P$.

Step 3. Q is Π_1^0 and recursively bounded. Hence, we can find Π_1^0 $Q^* \subseteq 2^\omega$ such that Q^* is recursively homeomorphic to Q . Done.

Some additional, specific degrees in \mathcal{P}_w :

\mathbf{d}_α = the weak degree of the set of diagonally nonrecursive functions which are bounded by a recursive function at level α of the Wainer hierarchy.

\mathbf{d}^2 = the weak degree of the set of $f \oplus g$ such that f is diagonally nonrecursive, and g is diagonally nonrecursive relative to f . More generally, define \mathbf{d}^n for all $n \geq 1$.

Theorem (Simpson 2004, Ambos et al 2004):
In \mathcal{P}_w we have

$$\mathbf{r}_1 > \mathbf{d}_0 > \mathbf{d}_1 > \cdots > \mathbf{d}_\alpha > \cdots > \mathbf{d}_{\text{REC}}$$

and

$$\mathbf{d} = \mathbf{d}^1 < \mathbf{d}^2 < \cdots < \mathbf{d}^n < \cdots < \mathbf{r}_1 .$$

We conjecture that \mathbf{d}^n is incomparable with \mathbf{d}_α and with \mathbf{d}_{REC} . This would be the first example of specific, natural degrees in \mathcal{P}_w which are incomparable with each other.

Response to Issue 2:

Issue 2 was:

To find a “smallness property” of an infinite Π_1^0 (i.e., co-r.e.) set $A \subseteq \omega$ which ensures that the Turing degree of A is $> \mathbf{0}$ and $< \mathbf{0}'$.

We do not know how to do this.

However, in the \mathcal{P}_ω context, we have identified several “smallness properties” of a Π_1^0 set $P \subseteq 2^\omega$ which ensure that the weak degree of P is $> \mathbf{0}$ and $< \mathbf{0}'$.

One result of this type:

Theorem (Simpson 2002):

Let \mathbf{p} be the weak degree of a Π_1^0 set $P \subseteq 2^\omega$ which is thin and perfect. Then \mathbf{p} is incomparable with \mathbf{r}_1 . Hence $\mathbf{0} < \mathbf{p} < \mathbf{0}'$.

Background on thin Π_1^0 sets:

Definition:

A Π_1^0 set $P \subseteq 2^\omega$ is said to be *thin* if, for all Π_1^0 sets $Q \subseteq P$, $P \setminus Q$ is Π_1^0 .

Thin perfect Π_1^0 subsets of 2^ω have been constructed by means of priority arguments. Much is known about them. For example, any two such sets are automorphic in the lattice of Π_1^0 subsets of 2^ω under inclusion.

(Martin/Pour-El 1970,
Downey/Jockusch/Stob 1990, 1996,
Cholak et al 2001)

Some additional “smallness properties” :

Let P be a Π_1^0 subset of 2^ω .

1. P is *small* if there is no recursive function f such that for all n there exist n members of P which differ at level $f(n)$ in the binary tree. (Binns 2003)

Example: Let $A \subseteq \omega$ be hypersimple, and let $A = B_1 \cup B_2$ where B_1, B_2 are r.e. Then $P = \{X \in 2^\omega \mid X \text{ separates } B_1, B_2\}$ is small.

Theorem (Binns):

If P is small, the weak degree of P is $< \mathbf{0}'$.

2. P is *h-small* if there is no recursive, canonically indexed sequence of pairwise disjoint clopen sets D_n , $n \in \omega$, such that $P \cap D_n \neq \emptyset$ for all n . (Simpson 2003)

Theorem (Simpson):

If P is h-small, the weak degree of P is $< \mathbf{0}'$.

Summary of this talk:

There are basic, unresolved issues concerning \mathcal{R}_T , the semilattice of recursively enumerable Turing degrees. One of the issues is the lack of specific, natural, r.e. degrees.

We embed \mathcal{R}_T into \mathcal{P}_w , the lattice of weak degrees of nonempty Π_1^0 subsets of 2^ω . We identify \mathcal{R}_T with its image in \mathcal{P}_w .

In the \mathcal{P}_w context, some of the unresolved issues can be satisfactorily addressed.

In particular, \mathcal{P}_w contains many specific, natural degrees which are related to foundationally interesting topics:

- algorithmic randomness,
- reverse mathematics,
- computational complexity.

References:

Binns, Small Π_1^0 classes, 22 pages,
Archive for Mathematical Logic, to appear.

Simpson, Mass problems and randomness,
Bulletin of Symbolic Logic, 11, 2005, 1–27.

Simpson, An extension of the recursively
enumerable Turing degrees, 15 pages,
submitted for publication.

Some of my papers are available at
<http://www.math.psu.edu/simpson/papers/>.

Transparencies for my talks are available at
<http://www.math.psu.edu/simpson/talks/>.

THE END