

# Operational measures for squeezing

**Martin Idel**

*joined work with **Daniel Lercher** and **Michael M. Wolf***

TNG Technology Consulting

24 July 2017



# What about the weird affiliation?



- Worked with Michael M. Wolf as a graduate student at TUM.
- Graduated recently and left academia.
- Work for TNG Technology Consulting:
  - Munich based German speaking IT development and consulting firm
  - > 50% PhD in Physics, Mathematics and Computer Science
  - special needs software consulting/development for various areas from telecommunications to autonomous driving
  - even a few quantum information people

# We pave the way to investigate squeezing as a resource

*If one specifies an error tolerance no larger than some error  $\varepsilon > 0$  and allows for using  $n$  instances of a given resource, what communication rates are achievable?*

In this talk:

- “New” resource theory with the usual questions: Squeezing of formation, distillation of squeezing, etc.
- Interesting due to connections with entanglement theory, experimental difficulties, maybe even on its own.
- Providing new tools to study the question in continuous variable quantum information.

# Modeling the electromagnetic field in phase space

- The electromagnetic field can be modeled as non-interacting harmonic oscillators (second quantisation).
- Harmonic oscillator description: frequency  $\omega_k$  and a set of position and momentum operators  $Q, P$ .
- Usually finitely many  $k$  are enough (e.g. in a cavity)  $\Rightarrow R = (Q_1, P_1, \dots, Q_n, P_n)$ .
- Photons are bosons  $\Rightarrow R$  fulfil the CCR:

$$[R_k, R_l] = i\sigma_{kl}\mathbb{1}, \quad \sigma = \bigoplus_{i=1}^n \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

- Symplectic transformations  $Sp(2n)$  leave the CCR invariant (corresponds to unitary transformations on the state).

# For Gaussian states, the covariance matrix is your friend

- $Q, P$  must be unbounded  $\Rightarrow$  use bounded representations  
 $W_\xi = \exp(-i\xi\sigma R)$ .
- Define the *characteristic function*  $\chi(\xi) = \text{tr}(W_\xi\rho)$ .
- The characteristic function of a state can often be described by its *moments*. **Gaussian states** are described by their first and second moments only:

$$d_k := \text{tr}(\rho R_k)$$

$$\gamma_{kl} := \text{tr}(\rho\{R_k - d_k\mathbb{1}, R_l - d_l\mathbb{1}\}_+).$$

- Operations on the state (such as time evolution) correspond to operations on the moments (such as symplectic transformations).
- **Heisenberg's uncertainty relation**:  $\gamma \geq i\sigma$ .
- A **squeezed state** has an eigenvalue  $\lambda$  of  $\gamma$  with  $\lambda < 1$ .

# Squeezing is a resource

- First noted by Braunstein: Squeezing remains invariant under linear optics [S.L. Braunstein. *PRA*, 71, 2005]
- **Free states:** one-mode squeezed states:  $\text{diag}(s, s^{-1})$
- free operations:
  1. Linear optics (symplectic orthogonal matrices  $S$  acting via  $\gamma \mapsto S^T \gamma S$ ),
  2. Free ancillary states ( $\gamma \rightarrow \gamma \oplus \gamma_{\text{anc}}$ ),
  3. Add classical noise ( $\gamma_{\text{noise}} \geq 0$  acting via  $\gamma \mapsto \gamma + \gamma_{\text{noise}}$ ),
  4. Weyl rotations (no change in covariance matrix),
  5. Convex combinations ( $\lambda\gamma + (1 - \lambda)\tilde{\gamma}$ ),
  6. Homodyne detection:

$$\gamma = \begin{pmatrix} A & C \\ C^T & B \end{pmatrix} \Rightarrow \gamma \mapsto \lim_{d \rightarrow \infty} A - C(B + \text{diag}(d, 1/d))^{-1} C^T.$$

# If you want entanglement, you need squeezing

## Theorem

Given a quantum state  $\rho$  with covariance matrix  $\gamma$ , for an arbitrary two-mode subsystem of a quantum state we have

$$E_N \propto \max\{0, -\log_2(\lambda_1 \lambda_2)\},$$

where  $\lambda_1, \lambda_2$  are the smallest eigenvalues of  $\gamma$  and  $E_N$  is the logarithmic negativity (an entanglement measure). [M.M. Wolf, J. Eisert, M. Plenio. PRL, 90, 2003]

## Theorem (No super-activation without squeezing)

Let  $T_1, T_2$  be passive Gaussian quantum channels. If each channel either has a symmetric extension or satisfies the PPT property, then

$$Q(T_1 \otimes T_2) = 0. \text{ [D. Lercher, G. Giedke, M.M. Wolf. New J. Phys. 15, 2013]}$$

# Squeezing measures as entanglement witnesses

- Variances (in position and momentum) can be measured very well in the lab.
- “Spin squeezing” measures have been used as entanglement measures for years.
- Similar “squeezing measures” have been proposed recently [M. Gessner, et al. *Quantum*, 2017-07-10]:

$$\xi^2(\gamma) = \min_{g \in \mathbb{R}^{2n}, \|g\|_2=1} (g^T \sigma^T \gamma_{\Pi(\rho)} \sigma g)(g^T \gamma_\rho g)$$

where  $\Pi(\rho) = \prod_{i=1}^N \rho_i$  with the reduced density matrices  $\rho_i$ . The separability criterion reads:

$$\xi^2(\gamma_{\text{sep}}) \geq 1$$

- Different goal: find entanglement, not study squeezing as is



# Current squeezing measures work well for one-mode states

Currently:  $G_{\text{squeeze}} = \lambda_{\min}(\gamma)$ . [B. Kraus et al. *PRA* 67:0402314, 2003]

Problems:

- Consider multimode states

$$\begin{pmatrix} s & & & \\ & \frac{1}{s} & & \\ & & 1 & \\ & & & 1 \end{pmatrix}, \begin{pmatrix} s & & & \\ & \frac{1}{s} & & \\ & & s & \\ & & & \frac{1}{s} \end{pmatrix}, \begin{pmatrix} s^2 & & & \\ & \frac{1}{s^2} & & \\ & & 1 & \\ & & & 1 \end{pmatrix}.$$

The first and second should not have the same squeezing.

- Operational measures?

## Defining measures of squeezing which also work for multiple modes

**Cost:** A one-mode squeezer  $S = \text{diag}(s, s^{-1})$  has costs  $\log(s)$ .

**Idea:** For a symplectic matrix  $S$  minimise one-mode squeezing for decompositions  $S = S_1 \dots S_m$  with  $S_i$  passive ( $K(n)$ ) or a one-mode squeezer ( $Z(n)$ ).

Then we have the measure:

$$F(S) = \inf \left\{ \sum_{i=1}^m \log s_1^\downarrow(S_i) \mid S = S_1 \dots S_m, S_i \in K(n) \cup Z(n) \right\}.$$

# Minimal squeezing for symplectic matrices is given by the Euler decomposition

Proposition (M.I., D. Lercher, M.M. Wolf (2016))

*For any symplectic matrix  $S$  we have*

$$F(S) = \sum_{i=1}^n \log(s_i^\downarrow(S)).$$

*Equality is achieved by the Euler decomposition*

*$S = K_1 \operatorname{diag}(s_1, s_1^{-1}, \dots, s_n, s_n^{-1}) K_2$  with passive  $K_1, K_2$ .*

## Proof

See Whiteboard

## This already is an operational measure for pure states

- If  $\text{diag}(s, s^{-1})$  are resource states, the optimal way to prepare a pure state with covariance matrix  $\gamma$  is given by the Euler decomposition.
- Preparation costs can be read from the Euler decomposition.
- For general states: Take a pure state and add noise.

Idea: Suggestion for a measure for general states:

$$G(\gamma) := \inf\{F(S) \mid \gamma \geq S^T S, S \in Sp(2n)\}$$

# Minimal squeezing for states is given by a simple optimisation problem

Minimise costs over all sequences

$$\gamma_0 \rightarrow \gamma_1 \rightarrow \dots \rightarrow \gamma_N = \gamma$$

with  $\gamma_0$  resource states and each operation being an allowed operation.

# Minimal squeezing for states is given by a simple optimisation problem

Minimise costs over all sequences

$$\gamma_0 \rightarrow \gamma_1 \rightarrow \dots \rightarrow \gamma_N = \gamma$$

with  $\gamma_0$  resource states and each operation being an allowed operation.

**Theorem (M.I., D. Lercher, M.M. Wolf (2016))**

*Given a quantum state  $\rho$  with covariance matrix  $\gamma$ , the minimal amount of one-mode squeezing needed for its creation is given by*

$$G(\gamma) = \min\{F(S) \mid \gamma \geq S^T S, S \in \text{Sp}(2n)\}$$

# Proof

See Whiteboard



# Detour: Cayley transformation

**Regular Cayley transform:** Transform upper complex half plane to unit disk.

**Matrix Cayley transform:**

- Transformation of positive half-plane  $Z > 0$  to unit disc  $\|H\| < 1$  [D. McDuff, D. Salamon, *Introduction to Symplectic Topology*, 1998].
- Transformation of skew-Hermitian matrix to unitary matrices.
- Transformation of symplectic matrices into Hamiltonian matrices [V. Mehrmann *Lin. Alg. App.* 241-243, 1996].
- Symplectic Cayley transform transforms symplectic matrices into symmetric ones [M. deGosson, *Symplectic Geometry and Quantum Mechanics*, 2006].

# Detour: Cayley transformation

$$C : \{H \in \mathbb{R}^{n \times n} \mid \text{spec}(H) \cap \{+1\} = \emptyset\} \mapsto \mathbb{R}^{n \times n}, \quad H \mapsto \frac{\mathbb{1} + H}{\mathbb{1} - H}.$$

$$C^{-1} : \{S \in \mathbb{R}^{n \times n} \mid \text{spec}(H) \cap \{-1\} = \emptyset\} \mapsto \mathbb{R}^{n \times n}, \quad S \mapsto \frac{S - \mathbb{1}}{S + \mathbb{1}}.$$

- $C$  and  $C^{-1}$  are diffeomorphisms onto their respective images.
- $C$  is operator monotone and operator convex on matrices  $A$  with  $\text{spec}(A) \subset (-1, 1)$ .
- $C^{-1}$  is operator monotone and operator concave on matrices  $A$  with  $\text{spec}(A) \subset (-1, \infty)$ .
- $C : \mathbb{R} \rightarrow \mathbb{R}$  is log-convex on  $[0, 1)$ .
- $C(\mathcal{H}) = \text{Sp}(2n) \cap \{\gamma \geq i\sigma\}$ , where

$$\mathcal{H} := \left\{ H = \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \middle| A \in \mathbb{R}^{2n \times 2n}, A^T = A, B^T = B, -\mathbb{1} < H < \mathbb{1} \right\}.$$

# We can rewrite $G$ using the Cayley transform

$$\begin{aligned}
 G(\gamma) &= \inf\{F(S) \mid \gamma \geq S^T S, S \in Sp(2n)\} \\
 &= \inf\{F(\gamma_0^{1/2}) \mid \gamma \geq \gamma_0 \geq iJ\} \\
 &= \inf \left\{ \frac{1}{2} \sum_{i=1}^n \log \left( \frac{1 + s_i(A + iB)}{1 - s_i(A + iB)} \right) \mid C^{-1}(\gamma) \geq H, H \in \mathcal{H} \right\}
 \end{aligned}$$

with (again)

$$\mathcal{H} := \left\{ H = \begin{pmatrix} A & B \\ B & -A \end{pmatrix} \mid A \in \mathbb{R}^{2n \times 2n} A^T = A, B^T = B, -\mathbb{1} < H < \mathbb{1} \right\}.$$


## Proof

See Whiteboard

# The measure improves preparation procedures

We provide numerical calculations for examples (two-parameter family of states, code can be found at <https://github.com/Martin-Idel/operationalsqueezing>):

[L. Mišta, N. Korolkova. PRA 77:050302, 2008]



# $G$ is superadditive and probably subadditive

We have:

$$\frac{1}{2}(G(\gamma_A) + G(\gamma_B)) \leq G(\gamma_A \oplus \gamma_B) \leq G(\gamma_A) + G(\gamma_B).$$

**Conjecture:**  $G$  is subadditive

- Supported by numerical data.
- True at least if  $\gamma_A$  is pure.

# We have bounds for $G$ , but the upper bound is bad

Best bounds:

$$-\frac{1}{2} \sum_{\lambda_i^\downarrow(\gamma) < 1} \log(\lambda_i^\downarrow(\gamma)) \leq G(\gamma) \leq F(S)$$

where  $S$  is the symplectic matrix in Williamson's theorem  $\gamma = S^T D S$ .

- lower bound achieved, if the eigenvectors to eigenvalues  $< 1$  can be extended to an orthonormal symplectic basis.
- upper bound can be arbitrarily bad: Thermal state with  $\gamma = n\mathbb{1}$  and  $S = \text{diag}(\sqrt{N-1}, 1/\sqrt{N-1}, \dots) \in Sp(2n)$ .

## Detour: Set-valued analysis

- Work with functions with *sets* and not just points as values.
- Define continuity, norms, etc.

### Definition

Let  $X, Y \subseteq \mathbb{R}^{n \times m}$  and  $f : X \rightarrow 2^Y$  be a set-valued function. It is *upper semicontinuous* (*upper hemicontinuous*) at  $x_0 \in X$  if:

for all open neighbourhoods  $Q$  of  $f(x_0)$  there exists an open neighbourhood  $W$  of  $x_0$  such that  $W \subseteq \{x \in X \mid f(x) \subset Q\}$ .

Likewise, we call it *lower semicontinuous* (often called *lower hemicontinuous*) at a point  $x_0$  if for any open set  $V$  intersecting  $f(x_0)$ , we can find a neighbourhood  $U$  of  $x_0$  such that  $f(x) \cap V \neq \emptyset$  for all  $x \in U$ .



## Detour: Set-valued analysis

Just for fun:

### Theorem

*Let  $S$  be a non-empty, compact and convex subset of some Euclidean space  $\mathbb{R}^n$ . Let  $f : S \rightarrow 2^S$  be a set-valued function on  $S$  with a closed graph and the property that  $f(x)$  is non-empty and convex for all  $x \in S$ . Then  $f$  has a fixed point.*

Proved in 1941 by Shizuo Kakutani and used in the one-page paper “Equilibrium points in  $N$ -person games” by John F. Nash.

# $G$ is probably continuous


Recall:

$$G(\gamma) = \inf\{F(\tilde{\gamma}^{1/2}) \mid \gamma \geq \tilde{\gamma} \geq i\sigma\}$$

For continuity, this means we have the intersection of two convex, non-empty and set-valued functions:

- $f(A) : A \geq i\sigma$ .
- $g(B) : B \leq \gamma$  (this one varies continuously).

**Heuristic:** This should be continuous.



# We should be able to prove continuity using set-valued analysis

**Current state:**  $G$  is lower semicontinuous on the set of covariance matrices and continuous on its interior.

**Conjecture:**  $G$  is continuous, since any compact intersection of set-valued functions consisting of matrix cones is continuous.

- Any intersection of non-empty compact sets with non-empty interior is continuous.
- Non-polynomial matrix cones make it somewhat difficult.

## Potential other applications

Would directly prove continuity of all functions consisting of optimisation of continuous functions over convex cones.

**Example:** Gaussian entanglement of formation: e.g. [M.M. Wolf., *PRA* **69**, 2004]

$$E_{\text{form}}(\gamma_{AB}) = \min\{H(\gamma_p) \mid \gamma_{AB} \geq S^T S, S \in Sp(2n)\}$$

where  $\gamma_p$  is the reduced state of  $S^T S$  and  $H$  is entanglement entropy.

# On the road to more realistic measures

- $\log(s)$  could be interpreted as “interaction time” of the squeezing Hamiltonian.
- In experiments: linear difficulty until cutoff at about 15 dB squeezing  
e.g. [H. Vahlbruch et al., *PRL* 117, 2016].
- Maybe exponential difficulty without explicit cutoff.

Problematic parts:

- Convexity of  $G$ .
- Submultiplicativity of  $F$  (irrelevant for resource theory).

## Can we do better using Lie algebras?

**Idea:** Maybe simple products of the form  $S = S_1 \cdots S_n$  are not optimal. How about general paths on the symplectic group?

**Proposition (M.I., D. Lercher, M.M. Wolf (2016))**

*General paths on  $Sp(2n)$  cannot decrease squeezing costs.*

We define the measure as follows:

$$\tilde{F}(S) := \inf \left\{ \int_0^1 \|\vec{c}_\alpha^a(t)\|_1 dt \mid \alpha \in C^r(S), \right. \\ \left. \dot{\alpha}(t) = (\vec{c}_\alpha^p(t)g^p(\alpha(t)), \vec{c}_\alpha^a(t)g^a(\alpha(t)))^T \right\}$$

with  $c_\alpha \in L^\infty([0, 1], \mathfrak{sp}(2n))$ .

# Proof

See Whiteboard

# Can we distill squeezing and use it in channels?

**Question:** What is the “Maximum output squeezing”. How to calculate it?  
Can the normal form for channels with squeezed environment help? [M.I., R. König. *Quant. Inf. Comp.*, 2017].

**Partial Answer:** Distillation of squeezing is prohibited for Gaussian states [B. Kraus, et al. *PRA* 67, 2003], hence not for the usual measure. For our measure, there are clear upper bounds  $\Rightarrow$  not so interesting.



# Can we distill squeezing and use it in channels?

**Question:** What is the “Maximum output squeezing”. How to calculate it?  
Can the normal form for channels with squeezed environment help? [M.I., R. König. *Quant. Inf. Comp.*, 2017].

**Partial Answer:** Distillation of squeezing is prohibited for Gaussian states [B. Kraus, et al. *PRA* 67, 2003], hence not for the usual measure. For our measure, there are clear upper bounds  $\Rightarrow$  not so interesting.

**Question:** What about non-gaussian states?  
Distillation is possible via linear optics and being studied [R. Filip. *PRA* 88, 2013].  
Problem: You cannot work with the covariance matrices only.

# I still have many open questions

- Fermionic quantum systems?
- Squeezing is related to the spectrum of the covariance matrix, while entanglement is related to the symplectic spectrum of submatrices. Can we have more explicit direct bounds?
- State interconvertibility is more complicated. Can we have even “better” measures?
- Can we have trade-off functions between squeezing and (e.g.) superactivation?