

Coding in graphs and linear orderings

Higher Recursion Theory and Set Theory

Alexandra A. Soskova ¹

Joint work with [J. Knight](#) and [S. Vatev](#)

¹Supported by Bulgarian National Science Fund DN 02/16 /19.12.2016 and NSF grant DMS 1600625/2016

Coding and decoding

- There are familiar ways of coding one structure in another, and for coding members of one class of structures in those of another class.
- Sometimes the coding is effective.
- Assuming this, it is interesting when there is effective decoding, and and it is also interesting when decoding is very difficult.

We consider some formal notions that describe coding and decoding, and test the notions in some examples.

Conventions

- 1 Languages are computable.
- 2 Structures have universe ω .
- 3 We may identify the structure \mathcal{A} with $D(\mathcal{A})$.
- 4 Classes \mathcal{K} are closed under isomorphism.

Borel embedding

Definition (Friedman, Stanley, 1989)

We say that a class \mathcal{K} of structures is *Borel embeddable* in a class of structures \mathcal{K}' , and we write $\mathcal{K} \leq_B \mathcal{K}'$, if there is a Borel function $\Phi : \mathcal{K} \rightarrow \mathcal{K}'$ such that for $\mathcal{A}, \mathcal{B} \in \mathcal{K}$, $\mathcal{A} \cong \mathcal{B}$ iff $\Phi(\mathcal{A}) \cong \Phi(\mathcal{B})$.

Note: We have a uniform Borel procedure for coding structures from structures of class \mathcal{K} in structures from \mathcal{K}' . As we shall see, there may or may not be a Borel decoding procedure.

On top

Theorem

The following classes lie on top under \leq_B .

- 1 undirected graphs (Lavrov, 1963; Nies, 1996; Marker, 2002)
- 2 fields of any fixed characteristic (Friedman-Stanley; R. Miller-Poonen-Schoutens-Shlapentokh, 2018)
- 3 2-step nilpotent groups (Mekler, 1981; Mal'tsev, 1949)
- 4 linear orderings (Friedman-Stanley)

Graphs \leq_B ACF(0)

Example

Let F^* be an algebraically closed field with transcendence basis b_0, b_1, b_2, \dots

For a graph G , let $F(G)$ be the subfield generated by the following:

- 1 b_i , for $i \in G$,
- 2 elements of $\text{acl}(b_i)$,
- 3 $\sqrt{d + d'}$, where for some i, j joined by an edge in G , d is inter-algebraic with b_i and d' is inter-algebraic with b_j .

The formulas that define the interpretation are computable Π_2^0 or simpler. Hence, for any $F \cong F(G)$, we get a copy of G computable in F'' .

Turing computable embeddings

Definition (Calvert, Cummins, Knight, S. Miller, 2004)

We say that a class \mathcal{K} is *Turing computably embedded* in a class \mathcal{K}' , and we write $\mathcal{K} \leq_{tc} \mathcal{K}'$, if there is a Turing operator $\Phi : K \rightarrow K'$ such that for all $\mathcal{A}, \mathcal{B} \in K$, $\mathcal{A} \cong \mathcal{B}$ iff $\Phi(\mathcal{A}) \cong \Phi(\mathcal{B})$.

The notion of Turing computable embedding captures in a precise way the idea of uniform effective coding.

On top

Theorem

The following classes lie on top under \leq_{tc} .

- 1 undirected graphs
- 2 fields of any fixed characteristic
- 3 2-step nilpotent groups
- 4 linear orderings

The Borel embeddings of Friedman and Stanley, Miller, Poonen, Schoutens and Shlapentokh, Lavrov, Nies, Marker, Mekler, and Mal'tsev are all, in fact, Turing computable.

Directed graphs \leq_{tc} undirected graphs

Example (Marker)

For a directed graph G the undirected graph $\Theta(G)$ consists of the following:

- 1 For each point a in G , $\Theta(G)$ has a point b_a connected to a triangle.
- 2 For each ordered pair of points $(a; a')$ from G , $\Theta(G)$ has a special point $p_{(a,a')}$ connected directly to b_a and with one stop to b'_a .
- 3 The point $p_{(a,a')}$ is connected to a square if there is an arrow from a to a' , and to a pentagon otherwise.

For structures \mathcal{A} with more relations, the same idea works.

Medvedev reducibility

A *problem* is a subset of 2^ω or ω^ω .

Problem P is Medvedev reducible to problem Q if there is a Turing operator Φ that takes elements of Q to elements of P .

Definition

We say that \mathcal{A} is *Medvedev reducible* to \mathcal{B} , and we write $\mathcal{A} \leq_s \mathcal{B}$, if there is a Turing operator that takes copies of \mathcal{B} to copies of \mathcal{A} .

Supposing that \mathcal{A} is coded in \mathcal{B} , a Medvedev reduction of \mathcal{A} to \mathcal{B} represents an effective decoding procedure.

For classes K and K' , suppose that $K \leq_{tc} K'$ via Θ . A uniform effective decoding procedure is a Turing operator Φ s.t. for all $\mathcal{A} \in K$, Φ takes copies of $\Theta(\mathcal{A})$ to copies of \mathcal{A} .

Decoding via nice defining formulas

Fact: For Marker's embedding Θ , we have finitary existential formulas that, for all directed graphs G , define in $\Theta(G)$ the following.

- 1 the set of points b_a connected to a triangle,
- 2 the set of ordered pairs such that the special point $p_{(a,a')}$ is part of a square,
- 3 the set of ordered pairs $(b_a, b_{a'})$ such that the special point $p_{(a,a')}$ is part of a pentagon.

This guarantees a uniform effective procedure that, for any copy of $\Theta(G)$, computes a copy of G . We have uniform effective decoding.

Effective interpretability

Definition (Harrison-Trainor, Melnikov, R. Miller, Montlbán)

A structure $\mathcal{A} = (A, R_i)$ is *effectively interpreted* in a structure \mathcal{B} if there is a set $D \subseteq \mathcal{B}^{<\omega}$ and relations \sim and R_i^* on D , such that

- 1 $(D, R_i^*)/\sim \cong \mathcal{A}$,
- 2 there are computable Σ_1 -formulas with no parameters defining a set $D \subseteq \mathcal{B}^{<\omega}$ and relations $(\neg)\sim$ and $(\neg)R_i^*$ in \mathcal{B} (effectively determined).

Example

The usual definition of the ring of integers \mathbb{Z} involves an interpretation in the semi-ring of natural numbers \mathbb{N} . Let D be the set of ordered pairs (m, n) of natural numbers. We think of the pair (m, n) as representing the integer $m - n$. We can easily give finitary existential formulas that define ternary relations of addition and multiplication on D , and the complements of these relations, and a congruence relation \sim on D , and the complement of this relation, such that $(D, +, \cdot)/\sim \cong \mathbb{Z}$.

Computable functor

Definition (Harrison-Trainor, Melnikov, R. Miller and Montalbán)

A *computable functor* from \mathcal{B} to \mathcal{A} is a pair of Turing operators Φ, Ψ such that Φ takes copies of \mathcal{B} to copies of \mathcal{A} and Ψ takes isomorphisms between copies of \mathcal{B} to isomorphisms between the corresponding copies of \mathcal{A} , so as to preserve identity and composition.

More precisely, Ψ is defined on triples $(\mathcal{B}_1, f, \mathcal{B}_2)$, where $\mathcal{B}_1, \mathcal{B}_2$ are copies of \mathcal{B} with $\mathcal{B}_1 \cong_f \mathcal{B}_2$.

Equivalence

The main result gives the equivalence of the two definitions.

Theorem (H-TMMM, 2017)

For structures \mathcal{A} and \mathcal{B} , \mathcal{A} is effectively interpreted in \mathcal{B} iff there is a computable functor Φ, Ψ from \mathcal{B} to \mathcal{A} .

Note: In the proof, it is important that D consist of tuples of arbitrary arity.

Corollary

If \mathcal{A} is effectively interpreted in \mathcal{B} , then $\mathcal{A} \leq_s \mathcal{B}$.

Coding and Decoding

Proposition (Kalimullin, 2010)

There exist \mathcal{A} and \mathcal{B} such that $\mathcal{A} \leq_s \mathcal{B}$ but \mathcal{A} is not effectively interpreted in \mathcal{B} .

There exist \mathcal{A} and \mathcal{B} such that \mathcal{A} is effectively interpreted in (\mathcal{B}, \bar{b}) but \mathcal{A} is not effectively interpreted in \mathcal{B} .

Proposition

If \mathcal{A} is computable, then it is effectively interpreted in all structures \mathcal{B} .

Proof.

Let $D = \mathcal{B}^{<\omega}$. Let $\bar{b} \sim \bar{c}$ if \bar{b}, \bar{c} are tuples of the same length. For simplicity, suppose $\mathcal{A} = (\omega, R)$, where R is binary. If $\mathcal{A} \models R(m, n)$, then $R^*(\bar{b}, \bar{c})$ for all \bar{b} of length m and \bar{c} of length n . Thus, $(D, R^*)/\sim \cong \mathcal{A}$. □

Parameters needed

Proposition

Suppose $\mathcal{B} \leq_s \mathcal{A}$. Then uniformly in a notation for α , the Σ_α^c -theory of \mathcal{B} is enumeration reducible to the Σ_α^c -theory of \mathcal{A} .

We have $\{\mathcal{A}\} \leq_{tc} \{\mathcal{B}\}$.

Example (Knight)

Let $\mathcal{B} = \mathcal{B}_1 + \{b\} + \mathcal{B}_2$, where, for a set S that is not Σ_3^0 , $\mathcal{B}_1 = \sigma(S \cup \{\omega\})$, and \mathcal{B}_2 is the shuffle sum of $\omega \cup \{\omega\}$. Without loss, we suppose that $1 \in S$. The orderings \mathcal{B} and \mathcal{B}_2 satisfy the same Σ_3 sentences— $\mathcal{B} \leq_3 \mathcal{B}_2$ and $\mathcal{B}_2 \leq_3 \mathcal{B}$. From an enumeration of the computable (or even finitary) Σ_3 formulas true of b , we can enumerate the set S . Since S is not Σ_3^0 , it is not enumeration reducible to the Σ_3^c theory of \mathcal{B}_2 , so it is not enumeration reducible to the Σ_3^c theory of \mathcal{B} . Therefore, $(\mathcal{B}, b) \not\leq_s \mathcal{B}$.

Borel interpretability

Harrison-Trainor, Miller and Montalbán, 2018, defined Borel versions of the notion of effective interpretation and computable functor.

Definition

- 1 For a Borel interpretation of $\mathcal{A} = (A, R_i)$ in \mathcal{B} the set $D \subseteq \mathcal{B}^{<\omega}$ the relations \sim and R_i^* on D , are definable by formulas of $L_{\omega_1\omega}$.
- 2 For a Borel functor from \mathcal{B} to \mathcal{A} , the operators Φ and Ψ are Borel.

Their main result gives the equivalence of the two definitions.

Theorem

A structure \mathcal{A} is interpreted in \mathcal{B} using $L_{\omega_1\omega}$ -formulas iff there is a Borel functor Φ, Ψ from \mathcal{B} to \mathcal{A} .

Graphs and linear orderings

Graphs and linear orderings both lie on top under Turing computable embeddings.

Graphs also lie on top under effective interpretation.

Question: What about linear orderings under effective interpretation?

And under using $L_{\omega_1\omega}$ -formulas?

Interpreting graphs in linear orderings

Proposition (Knight-S.-Vatev)

There is a graph G such that for all linear orderings L , $G \not\leq_s L$.

Proof.

Let S be a non-computable set. Let G be a graph such that every copy computes S .

We may take G to be a “daisy” graph”, consisting of a center node with a “petal” of length $2n + 3$ if $n \in S$ and $2n + 4$ if $n \notin S$.

Now, apply:

Proposition (Richter)

For a linear ordering L , the only sets computable in all copies of L are the computable sets.



Interpreting a graph in the jump of linear ordering

We are identifying a structure \mathcal{A} with its atomic diagram. We may consider an interpretation of \mathcal{A} in the jump \mathcal{B}' of \mathcal{B} . Note that the relations definable in \mathcal{B}' by computable Σ_1 relations are the ones definable in \mathcal{B} by computable Σ_2 relations.

Proposition (Knight-S.-Vatev)

There is a graph G such that for all linear orderings L , $G \not\leq_S L'$.

Proof.

Let S be a non- Δ_2^0 set. Let G be a graph such that every copy computes S . Then apply:

Proposition (Knight, 1986)

For a linear ordering L , the only sets computable in all copies of L' (or in the jumps of all copies of L), are the Δ_2^0 sets.



Interpreting a graph in the second jump of linear ordering

Proposition

For any set S , there is a linear ordering L such that for all copies of L , the second jump computes S .

Proof.

We may take L to be a “shuffle sum” of $n + 1$ for $n \in S \oplus S^c$ and ω . \square

Proposition

For any graph G , there is a linear ordering L such that $G \leq_s L''$. In fact, G is interpreted in L using computable Σ_3 formulas.

Proof.

Let S be the diagram of a specific copy G_0 of G and let L be a linear order such that $S \leq_s L''$. We have computable functor that takes the second jump of any copy of L to G_0 , and takes all isomorphisms between copies of L to the identity isomorphism on G_0 . \square

Friedman-Stanley embedding of graphs in orderings

Friedman and Stanley determined a Turing computable embedding $L : G \rightarrow L(G)$, where $L(G)$ is a sub-ordering of $\mathbb{Q}^{<\omega}$ under the lexicographic ordering.

- 1 Let $(A_n)_{n \in \omega}$ be an effective partition of \mathbb{Q} into disjoint dense sets.
- 2 Let $(t_n)_{1 \leq n}$ be a list of the atomic types in the language of directed graphs.

Definition

For a graph G , the elements of $L(G)$ are the finite sequences $r_0 q_1 r_1 \dots r_{n-1} q_n r_n k \in \mathbb{Q}^{<\omega}$ such that for $i < n$, $r_i \in A_0$, $r_n \in A_1$, and for some $a_1, \dots, a_n \in G$, satisfying t_m , $q_i \in A_{a_i}$ and $k < m$.

Properties of $L(G)$

Definition

Let $b = r_0 q_1 r_1 \dots r_{n-1} q_n r_n k \in L(G)$, and let \bar{a} be the tuple in G such that $q_i \in A_{a_i}$. Then we say that b mentions \bar{a} .

Lemma

Suppose $b \in L(G)$ mentions \bar{a} . Then b lies in maximal discrete interval of some finite size $m \geq 1$.

Note that if b mentions \bar{a} of length n , then b has length $2n + 2$.

Lemma

If $b \in L(G)$ has length $2n + 2$, then there is an infinite interval around b that consists entirely of elements of length at least $2n + 2$.

Lemma

Let $b < b'$ in $L(G)$, and let d be an element of $[b, b']$ of minimum length. If d mentions \bar{c} , then all elements of $[b, b']$ mention extensions of \bar{c} .

No uniform interpretation of G in $L(G)$

Theorem (Knight-S.-Vatev, HarrisonTrainor-Montlbán)

There are no $L_{\omega_1\omega}$ formulas that, for all graphs G , interpret G in $L(G)$.

The idea of Proof by Knight-S.-Vatev: We may think of an ordering as a directed graph. It is enough to show the following.

Proposition

- A ω_1^{CK} is not interpreted in $L(\omega_1^{CK})$ using computable infinitary formulas.
- B For all X , ω_1^X is not interpreted in $L(\omega_1^X)$ using X -computable infinitary formulas.

Proof of A

The **Harrison ordering** H has order type $\omega_1^{CK}(1 + \eta)$. It has a computable copy.

Let I be the initial segment of H of order type ω_1^{CK} . Thinking of H as a directed graph, we can form the linear ordering $L(H)$. We consider $L(I) \subseteq L(H)$.

Lemma

$L(I)$ is a computable infinitary elementary substructure of $L(H)$.

Proposition (Main)

There do not exist computable infinitary formulas that define an interpretation of H in $L(H)$ and an interpretation of I in $L(I)$.

To prove A, we suppose that there are computable infinitary formulas interpreting ω_1^{CK} in $L(\omega_1^{CK})$. Using Barwise Compactness theorem, we get essentially H and I with these formulas interpreting H in $L(H)$ and I in $L(I)$.

Proof of the Proposition(Main)

Lemma

- 1 For any $\bar{b} \in L(I)$, and $c \in L(I)$ there is an automorphism of $L(I)$ taking \bar{b} to a tuple \bar{b}' entirely to the right of c .
- 2 For any $\bar{b} \in L(I)$, and $c \in L(I)$ there is also an automorphism taking \bar{b} to a tuple \bar{b}'' entirely to the left of c .

Lemma

Suppose that we have computable Σ_γ formulas D , \otimes and \sim , defining an interpretation of H in $L(H)$ and I in $L(I)$. Then in $D^{L(I)}$ there is a fixed n , and there are n -tuples, all satisfying the same Σ_γ formulas, and representing arbitrarily large ordinals $\alpha < \omega_1^{CK}$.

We arrive at a contradiction by producing tuples $\bar{b}, \bar{b}', \bar{c}$ in $D^{L(I)}$, \bar{b} and \bar{b}' are automorphic, \bar{b}, \bar{c} and \bar{c}, \bar{b}' satisfy the same Σ_γ formulas, and the ordinal represented by \bar{b} and \bar{b}' is smaller than that represented by \bar{c} . Then \bar{b}, \bar{c} should satisfy \otimes , while \bar{c}, \bar{b}' should not.

Conjecture

We believe that Friedman and Stanley did the best that could be done.

Conjecture. For any Turing computable embedding Θ of graphs in orderings, there do not exist $L_{\omega_1\omega}$ formulas that, for all graphs G , define an interpretation of G in $\Theta(G)$.

M. Harrison-Trainor and A. Montalbán came to a similar result recently by a totally different construction. Their result is that there exist structures which cannot be computably recovered from their tree of tuples. They proved :

- 1 There is a structure \mathcal{A} with no computable copy such that $T(\mathcal{A})$ has a computable copy.
- 2 For each computable ordinal α there is a structure \mathcal{A} such that the Friedman and Stanley Borel interpretation $L(\mathcal{A})$ is computable but \mathcal{A} has no Δ_α^0 copy.

Mal'tsev embedding of fields in groups

If F is a field, we denote by $H(F)$ the multiplicative group of matrices of kind

$$h(a, b, c) = \begin{pmatrix} 1 & a & b \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}$$

where $a, b, c \in F$. Note that $h(0, 0, 0) = 1$.

Groups of kind $H(F)$ are known as *Heisenberg groups*.

Theorem (Mal'tsev)

There is a copy of F defined in $H(F)$ with parameters.

Definition of F in $H(F)$

Let u, v be a non-commuting pair in $H(F)$.

Then $(D, +, \cdot_{(u,v)})$ is a copy of F , where

- 1 D is the group center – $x \in D \iff [x, u] = 1$ and $[x, v] = 1$,
- 2 $x + y = z$ if $x * y = z$, where $*$ is the group operation,
- 3 $x \cdot_{(u,v)} y = z$ if there exist x', y' such that $[x', u] = [y', v] = 1$, $[x', v] = x$, $[u, y'] = y$, and $[x', y'] = z$.

Here $[x, y] = x^{-1}y^{-1}xy$.

Definability: We have finitary existential formulas that define D and the relation $+$ and its complement. For any non-commuting pair (u, v) , we have finitary existential formulas, with parameters (u, v) that define the relation \cdot and its complement.

Natural isomorphisms

For a non-commuting pair (u, v) , where $u = h(u_1, u_2, u_3)$ and $v = h(v_1, v_2, v_3)$, let

$$\Delta_{(u,v)} = \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix}$$

Theorem (Morozov)

The function f that takes $x \in F$ to $h(0, 0, \Delta_{(u,v)} \cdot_F x)$ is an isomorphism.

Morozov's isomorphism

Lemma

Let (u, v) and (u', v') be non-commuting pairs in $G = H(F)$. Let $F_{(u,v)}$ and $F_{(u',v')}$ be the copies of F defined in G with these pairs of parameters. There is an isomorphism g from $F_{(u,v)}$ onto $F_{(u',v')}$ defined in G by an existential formula with parameters u, v, u', v' .

Note that $\Delta_{(u,v)}$ is the multiplicative identity in $F_{(u,v)}$.

Let $g(x) = y \iff x = \Delta_{(u,v)} \cdot (u',v') y$.

Computable functor

Theorem

There is a computable functor Φ, Ψ from $H(F)$ to F .

- For $G \cong H(F)$, $\Phi(G)$ is the copy of F obtained by taking the first non-commuting pair (u, v) in G and forming $(D; +; \cdot_{(u,v)})$.
- Take (G_1, f, G_2) , where $G_i = H(F)$, and $G_1 \cong_f G_2$. Let $(u, v), (u', v')$ be the first non-commuting pairs in G_1, G_2 , respectively.
 - ▶ Let h be the isomorphism from $F_{(f(u), f(v))}$ onto $F_{(u', v')}$ defined in G_2 with parameters $f(u), f(v), u', v'$.
 - ▶ Let f' be the restriction of f to the center of G_1 .
 - ▶ Then $\Psi(G_1, f, G_2) = h \circ f'$.

Finitely existential interpretation and generalizing

Corollary (Alvir, Calvert, Harizanov, Knight, Miller, Morozov, S, Weisshaar)

F is effectively interpreted in $H(F)$.

$(u, v, x) \sim (u', v', x')$ holds if Morozov's isomorphism from $F_{(u,v)}$ to $F_{(u',v')}$ takes x to x' .

Proposition

Suppose \mathcal{A} has a copy $\mathcal{A}_{\bar{b}}$ defined in (\mathcal{B}, \bar{b}) , using computable Σ_1 formulas, where the orbit of \bar{b} is defined by a computable Σ_1 formula $\varphi(\bar{x})$. Suppose also that there is a computable Σ_1 formula $\psi(\bar{b}, \bar{b}', u, v)$ that, for any tuples \bar{b}, \bar{b}' satisfying $\varphi(\bar{x})$, defines a specific isomorphism $f_{\bar{b}, \bar{b}'}$ from $\mathcal{A}_{\bar{b}}$ onto $\mathcal{A}_{\bar{b}'}$. We suppose that for each \bar{b} satisfying φ , $f_{\bar{b}, \bar{b}}$ is the identity isomorphism, and for any \bar{b}, \bar{b}' , and \bar{b}'' satisfying φ , $f_{\bar{b}', \bar{b}''} \circ f_{\bar{b}, \bar{b}'} = f_{\bar{b}, \bar{b}''}$. Then there is an effective interpretation of \mathcal{A} in \mathcal{B} .

$SL_2(C)$

Let C be an algebraically closed field of characteristic 0 and of infinite transcendence degree.

We consider $SL_2(C)$ for the group of 2×2 matrices over C with determinant 1.

Proposition

F is interpreted in $SL_2(F)$ with parameters.

Let A be the set of matrices of form $\begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$.

Let M be the set of matrices of form $\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$.

$SL_2(C)$

Let T consist of the pairs (X, Y) such that $X \in A$ and $Y \in M$ and Y has a square root Z such that $Z * P * Z^{-1} = X$.

For $(X, Y) \in T$, we define addition and multiplication relations as follows:

- 1 $(X, Y) \oplus (X', Y') = (U, V)$ if $X * X' = U$ and $(U, V) \in T$,
- 2 $(X, Y) \otimes (X', Y') = (U, V)$ if $Y * Y' = V$ and $(U, V) \in T$.

We define the set T with parameters.

Possibly, we can show model completeness of the theory of $SL_2(C)$.

This, together with the result, according to Pillay, saying that C is interpreted in $SL_2(C)$ by elementary first order formulas with no parameters, we could show that it is interpreted with existential formulas.



W. Calvert, D. Cummins, J. F. Knight, and S. Miller

Comparing classes of finite structures

Algebra and Logic, vo. 43(2004), pp. 374-392.



H. Friedman and L. Stanley

A Borel reducibility theory for classes of countable structures

JSL, vol. 54(1989), pp. 894-914.



J. Harrison,

Recursive pseudo well-orderings

Transactions of the Amer. Math. Soc., vol. 131 (1968) pp. 526-543.



M. Harrison-Trainor, A. Melnikov, R. Miller, and A. Montalbán

Computable functors and effective interpretability,

JSL, vol. 82(2017), pp. 77-97.



M. Harrison-Trainor, R. Miller, and A. Montalbán

Borel functors and infinitary interpretations,

JSL, vol. 83(2018), no. 4, pp. 1434-1456.

THANK YOU



Thank you for your deep contributions to logic!

Happy anniversary!